

09-05-00

A

Customer No. 000959

Case Docket No. UIZ-038

09/01/00

Jc836 U.S. PTO
09/653730
09/01/00

ASSISTANT COMMISSIONER FOR PATENTS
Box Patent Application
Washington, D.C. 20231

"Express Mail" Mailing Label Number EL 373209892 US

Date of Deposit September 1, 2000

I hereby certify that this transmittal letter and the papers referred to as being enclosed therein are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

Larry Taylor
Signature

Larry Taylor

Please Print Name of Person Signing

Sir:

Transmitted herewith for filing is the patent application of

Inventor(s): Marvin Whiteley, Kimberly M. Lee, E. Peter Greenberg, Ute Muh

For: QUORUM SENSING SIGNALING IN BACTERIA

Enclosed are:

- ☒ This is a request for filing an application under 37 CFR 1.53(b).
- ☒ 54 pages of specification, 10 pages of claims, 1 page of abstract.
- ☒ 13 sheets of informal drawings (Figures 1-9, 10A, 10B, 11, 12).
- ☒ An unexecuted Declaration, Petition and Power of Attorney.
- ☒ 24 pages of sequence listing (numbered 1- 24)
- ☒ Transmittal Letter for Diskette of Sequence Listing.
- ☒ Diskette Containing Sequence Listing.

The filing fee has been calculated as shown below:

	(Col. 1)	(Col. 2)
FOR:	NO. FILED	NO. EXTRA
BASIC FEE	////////////////////	
TOTAL CLAIMS	91-20	= 71
INDEP. CLAIMS	9 - 3	= 6
<input checked="" type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED		

* If the difference in Col. 2 is less than zero,
enter "0" in Col. 2.

SMALL ENTITY

RATE	FEE
////////	\$ 345
x 9=	\$ 639
x 39	\$ 234
+130	\$ 130
TOTAL	\$1348.00

OTHER THAN
SMALL ENTITY

OR	RATE	FEE
OR	////////	\$
OR	x 18=	\$
OR	x 78	\$
OR	+260	\$
OR	TOTAL	\$.00

- ☐ Please charge my Deposit Account No. in the amount of \$.
A duplicate copy of this sheet is enclosed.
- ☒ **FILING FEES ARE NOT BEING PAID AT THIS TIME.**
- ☐ The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No..
A duplicate copy of this sheet is enclosed.
- ☐ Any additional filing fees required under 37 C.F.R. 1.16.
- ☐ Any patent application processing fees under 37 C.F.R. 1.17.
- ☐ The Commissioner is hereby authorized to charge payment of the following fees during the pendency of this application or credit any overpayment to Deposit Account No..
A duplicate copy of this sheet is enclosed.

- Lahive & Cockfield, LLP
28 State Street
Boston, Massachusetts 02109

Date: September 1, 2000

By Peter C. Lauro
Peter C. Lauro, Esq.
Reg. No. 32,360
28 State Street
Boston, MA 02109
(617) 227-7400
Telecopier (617) 742-4214

Related Applications

Government Support

Background of the Invention

The paradigm system for quorum sensing is the *lux* system of the luminescent marine bacterium, *Vibrio fischeri*. *V. fischeri* exists at low cell densities in sea water and also at very high cell densities within the light organs of various marine organisms, such as the squid *Euprymna scolopes* (Pesci, E.C. *et al.* (1997) *Trends in Microbiol.* 5(4):132-135; Pesci, E.C. *et al.* (1997) *J. Bacteriol.* 179:3127-3132; Ruby, E.G. (1996) *Ann. Rev. Microbiol.* 50:591-624). At high cell densities, the *V. fischeri* genes encoding the enzymes required for light production are expressed. These genes are part of the *lux ICDABEG* operon and are regulated by the gene products of *luxI* and *luxR* (Baldwin, T.O. *et al.* (1989) *J. of Biolum. and Chemilum.* 4:326-341; Eberhard, A., *et al.* (1991) *Arch. of Microbiol.* 155:294-297; Gray, K.M. *et al.* (1992) *J. Bacteriol.* 174:4384-4390).

LuxI is an autoinducer synthase that catalyzes the formation of the *V. fischeri* autoinducer (VAI), *N*-(3-oxohexanoyl) homoserine lactone (Eberhard, A., *et al.* (1991) *Arch. of Microbiol.* 155:294-297; Seed, P.C. *et al.* (1995) *J. Bacteriol.* 177:654-659).

35 The autoinducer freely diffuses across the cell membrane and at high cell densities, reaches a critical concentration (Kaplan, H.B. *et al.* (1985) *J. Bacteriol.* 163:1210-1214). At this critical concentration, VAI interacts with LuxR, a DNA-binding transcriptional regulator. The LuxR-VAI complex then binds to an upstream sequence

of the *lux* operon called the "lux box", and activates transcription (Devine, J.H. *et al.* (1989) *PNAS* 86: 5688-5692; Hanzelka, B.A. *et al.* (1995) *J Bacteriol.* 177:815-817; Stevens, A.M. *et al.* (1994) *PNAS* 91:12619-12623). Since one of the genes of the operon is *luxI*, an autoregulatory loop is formed.

5 Many gram-negative bacteria have been shown to possess one or more quorum sensing systems (Fuqua, W.C. *et al.* (1996) *Annu. Rev. Microbiol.* 50:727-751; Salmond, G.P.C. *et al.* (1995) *Mol. Microbiol.* 16:615-624). These systems regulate a variety of physiological processes, including the activation of virulence genes and the formation of biofilms. The systems typically have acylated homoserine lactone ring
10 autoinducers, in which the homoserine lactone ring is conserved. The acyl side chain, however, can vary in length and degree of substitution. In addition, it has been recently demonstrated that quorum sensing is involved in biofilm formation (Davies, D. G. *et al.* (1998) *Science.* 280(5361):295-8).

Biofilms are defined as an association of microorganisms, single or multiple
15 species, that grow attached to a surface and produce a slime layer that provides a protective environment (Costerton, J. W. (1995) *J Ind Microbiol.* 15(3):137-40, Costerton, J. W. *et al.* (1995) *Annu Rev Microbiol.* 49:711-45). Typically, biofilms produce large amounts of extracellular polysaccharides, responsible for the slimy appearance, and are characterized by an increased resistance to antibiotics (1000- to
20 1500-fold less susceptible). Several mechanisms are proposed to explain this biofilm resistance to antimicrobial agents (Costerton, J. W. *et al.* (1999) *Science.* 284(5418):1318-22). One idea is that the extracellular matrix in which the bacterial cells are embedded provides a barrier toward penetration by the biocides. A further possibility is that a majority of the cells in a biofilm are in a slow-growing, nutrient-
25 starved state, and therefore not as susceptible to the effects of anti-microbial agents. A third mechanism of resistance could be that the cells in a biofilm adopt a distinct and protected biofilm phenotype, *e.g.*, by elevated expression of drug-efflux pumps.

In most natural settings, bacteria grow predominantly in biofilms. Biofilms of *P. aeruginosa* have been isolated from medical implants, such as indwelling urethral,
30 venous or peritoneal catheters (Stickler, D. J. *et al.* (1998) *Appl Environ Microbiol.* 64(9):3486-90). Chronic *P. aeruginosa* infections in cystic fibrosis lungs are considered to be biofilms (Costerton, J. W. *et al.* (1999) *Science.* 284(5418):1318-22).

In industrial settings, the formation of biofilms is often referred to as 'biofouling'. Biological fouling of surfaces is common and leads to material
35 degradation, product contamination, mechanical blockage, and impedance of heat transfer in water-processing systems. Biofilms are also the primary cause of biological contamination of drinking water distribution systems, due to growth on filtration devices.

007060-0229950

As noted earlier, many gram-negative bacteria have been shown to possess one or more quorum sensing systems that regulate a variety of physiological processes, including the activation of virulence genes and biofilm formation. One such gram negative bacterium is *Pseudomonas aeruginosa*.

5 *P. aeruginosa* is a soil and water bacterium that can infect animal hosts. Normally, the host defense system is adequate to prevent infection. However, in immunocompromised individuals (such as burn patients, patients with cystic fibrosis, or patients undergoing immunosuppressive therapy), *P. aeruginosa* is an opportunistic pathogen, and infection with *P. aeruginosa* can be fatal (Govan, J. R. *et al.* (1996) *Microbiol Rev.* 60(3):539-74; Van Delden, C. *et al.* (1998) *Emerg Infect Dis.* 4(4):551-60).

For example, Cystic fibrosis (CF), the most common inherited lethal disorder in Caucasian populations (~1 out of 2,500 life births), is characterized by bacterial colonization and chronic infections of the lungs. The most prominent bacterium in these
15 infections is *P. aeruginosa*—by their mid-twenties, over 80% of people with CF have *P. aeruginosa* in their lungs (Govan, J. R. *et al.* (1996) *Microbiol Rev.* 60(3):539-74). Although these infections can be controlled for many years by antibiotics, ultimately they “progress to mucoidy,” meaning that the *P. aeruginosa* forms a biofilm that is resistant to antibiotic treatment. At this point the prognosis is poor. The median
20 survival age for people with CF is the late 20s, with *P. aeruginosa* being the leading cause of death (Govan, J. R. *et al.* (1996) *Microbiol Rev.* 60(3):539-74). According to the Cystic Fibrosis Foundation, treatment of CF cost more than \$900 million in 1995 (Foundation, CF <http://www.cff.org/homeline199701.htm>).

P. aeruginosa is also one of several opportunistic pathogens that infect people
25 with AIDS, and is the main cause of bacteremia (bacterial infection of the blood) and pneumonitis in these patients (Rolston, K. V. *et al.* (1990) *Cancer Detect Prev.* 14(3):377-81; Witt, D. J. *et al.* (1987) *Am J Med.* 82(5):900-6). A recent study of 1635 AIDS patients admitted to a French hospital between 1991-1995 documented 41 cases of severe *P. aeruginosa* infection (Meynard, J. L. *et al.* (1999) *J Infect.* 38(3):176-81).
30 Seventeen of these had bacteremia, which was lethal in 8 cases. Similar, numbers were obtained in a smaller study in a New York hospital, where the mortality rate for AIDS patients admitted with *P. aeruginosa* bacteremia was about 50% (Mendelson, M. H. *et al.* 1994. *Clin Infect Dis.* 18(6):886-95).

In addition, about two million Americans suffer serious burns each year, and
35 10,000-12,000 die from their injuries. The leading cause of death is infection (Lee, J. J. *et al.* (1990) *J Burn Care Rehabil.* 11(6):575-80). *P. aeruginosa* bacteremia occurs in 10 % of seriously burned patients, with a mortality rate of 80 % (Mayhall, C. G. (1993) p.

004660 0646900

614-664, Prevention and control of nosocomial infections. Williams & Wilkins, Baltimore; McManus, A. T *et al.* (1985) *Eur J Clin Microbiol.* 4(2):219-23).

Such infections are often acquired in hospitals ("nosocomial infections") when susceptible patients come into contact with other patients, hospital staff, or equipment.

5 In 1995 there were approximately 2 million incidents of nosocomial infections in the U.S., resulting in 88,000 deaths and an estimated cost of \$ 4.5 billion (Weinstein, R. A. (1998) *Emerg Infect Dis.* 4(3):416-20). Of the AIDS patients mentioned above who died of *P. aeruginosa* bacteremia, more than half acquired these infections in hospitals (Meynard, J. L. *et al.* (1999) *J Infect.* 38(3):176-81).

10 Nosocomial infections are especially common in patients in intensive care units as these people often have weakened immune systems and are frequently on ventilators and/or catheters. Catheter-associated urinary tract infections are the most common nosocomial infection (Richards, M. J. *et al.* (1999) *Crit Care Med.* 27(5):887-92) (31 % of the total), and *P. aeruginosa* is highly associated with biofilm growth and catheter
15 obstruction. While the catheter is in place, these infections are difficult to eliminate (Stickler, D. J. *et al.* (1998) *Appl Environ Microbiol.* 64(9):3486-90). The second most frequent nosocomial infection is pneumonia, with *P. aeruginosa* the cause of infection in 21 % of the reported cases (Richards, M. J. *et al.* (1999) *Crit Care Med.* 27(5):887-92). The annual costs for diagnosing and treating nosocomial pneumonia has been
20 estimated at greater than \$2 billion (Craven, D. E. *et al.* (1991) *Am J Med.* 91(3B):44S-53S).

Treatment of these so-called nosocomial infections is complicated by the fact that bacteria encountered in hospital settings are often resistant to many antibiotics. In June 1998, the National Nosocomial Infections Surveillance (NNIS) System reported
25 increases in resistance of *P. aeruginosa* isolates from intensive care units of 89 % for quinolone resistance and 32 % for imipenem resistance compared to the years 1993-1997 (NNIS. http://www.cdc.gov/ncidod/hip/NNIS/AR_Surv1198.htm). In fact, some strains of *P. aeruginosa* are resistant to over 100 antibiotics (Levy, S. (1998) *Scientific American*, March). There is a critical need to overcome the emergence of bacterial
30 strains that are resistant to conventional antibiotics (Travis, J. (1994) *Science.* 264:360-362).

P. aeruginosa is also of great industrial concern (Bitton, G. (1994) *Wastewater Microbiology*. Wiley-Liss, New York, NY; Steelhammer, J. C. *et al.* (1995) *Indust. Water Treatm.* :49-55). The organism grows in an aggregated state, the biofilm, which
35 causes problems in many water processing plants. Of particular public health concern are food processing and water purification plants. Problems include corroded pipes, loss of efficiency in heat exchangers and cooling towers, plugged water injection jets leading to increased hydraulic pressure, and biological contamination of drinking water

U I Z - 0 3 8

A particularly ironic connection between industrial water contamination and public health issues is an outbreak of *P. aeruginosa* peritonitis that was traced back to contaminated poloxamer-iodine solution, a disinfectant used to treat the peritoneal catheters. *P. aeruginosa* is commonly found to contaminate distribution pipes and water filters used in plants that manufacture iodine solutions. Once the organism has matured into a biofilm, it becomes protected against the biocidal activity of the iodophor solution. Hence, a common soil organism that is harmless to the healthy population, but causes mechanical problems in industrial settings, ultimately contaminated antibacterial solutions that were used to treat the very people most susceptible to infection.

Summary of the Invention

35 Thus, in one aspect, the invention is a method for identifying a modulator of quorum sensing signaling in bacteria. The method comprises:

providing a cell comprising a quorum sensing controlled gene, wherein the cell is responsive to a quorum sensing signal molecule such that a detectable signal is generated;

5 contacting said cell with a quorum sensing signal molecule in the presence and absence of a test compound;

and detecting a change in the detectable signal to thereby identify the test compound as a modulator of quorum sensing signaling in bacteria.

10 In one embodiment the cell comprises a reporter gene operatively linked to a regulatory sequence of a quorum sensing controlled gene, such that the quorum sensing signal molecule modulates the transcription of the reporter gene, thereby providing a detectable signal.

Another aspect of the invention is a method for identifying a modulator of a quorum sensing signaling in *Pseudomonas aeruginosa*. The method comprises:

15 providing a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule;

providing a mutant strain of *Pseudomonas aeruginosa* which comprises a reporter gene operatively linked to a regulatory sequence of a quorum sensing controlled gene, wherein the mutant strain is responsive to the quorum sensing signal molecule produced by the wild type strain, such that a detectable signal is generated;

20 contacting the mutant strain with the quorum sensing signal molecule and a test compound; and

detecting a change in the detectable signal to thereby identify the test compound as a modulator of quorum sensing signaling in *Pseudomonas aeruginosa*.

25 In one embodiment, the endogenous *lasI* and *rhII* quorum sensing systems are inactivated in the mutant strain of *Pseudomonas aeruginosa*. In another embodiment the mutant strain of *Pseudomonas aeruginosa* comprises a promoterless reporter gene inserted at a genetic locus in the chromosome, wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID
30 NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID
35 NO:34, SEQ ID NO:35 and SEQ ID NO:36.

A further aspect of the invention is a mutant strain of *Pseudomonas aeruginosa* comprising a promoterless reporter gene inserted at a genetic locus in the chromosome, wherein the genetic locus comprises a nucleotide sequence selected from the group

004660086950

consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36.

In one embodiment, the endogenous *lasI* and *rhlI* quorum sensing systems are inactivated in the mutant strain of *Pseudomonas aeruginosa*. In another embodiment the mutant strain of *Pseudomonas aeruginosa* is responsive to a quorum sensing signal molecule such that a detectable signal is generated by the reporter gene. In yet another embodiment, the reporter gene is contained in a transposable element.

Yet another aspect of the invention is a method for identifying a modulator of quorum sensing signaling in *Pseudomonas aeruginosa*. The method comprises:

providing a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule;

providing a mutant strain of *Pseudomonas aeruginosa* which comprises a promoterless reporter gene inserted at a genetic locus in the chromosome of said *Pseudomonas aeruginosa*, wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36; and wherein the mutant strain is responsive to the quorum sensing signal molecule produced by the wild type strain, such that a detectable signal is generated by the reporter gene;

contacting the mutant strain with the quorum sensing signal molecule and a test compound; and

detecting a change in the detectable signal to thereby identify the test compound as a modulator of quorum sensing signaling in *Pseudomonas aeruginosa*.

Another aspect of the invention is an isolated nucleic acid molecule comprising a nucleotide sequence which comprises:

a regulatory sequence derived from the genome of *Pseudomonas aeruginosa*, wherein the regulatory sequence regulates a quorum sensing controlled genetic locus of

the *Pseudomonas aeruginosa* chromosome, and wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36; and

10 a reporter gene operatively linked to the regulatory sequence.

A further aspect of the invention provides an isolated nucleic acid molecule comprising a quorum sensing controlled genetic locus derived from the genome of *Pseudomonas aeruginosa*, wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36, operatively linked to a reporter gene.

In one embodiment, the invention is an isolated nucleic acid molecule comprising a polynucleotide having at least 80% identity to a quorum sensing controlled genetic locus derived from the genome of *Pseudomonas aeruginosa*, wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36, operatively linked to a reporter gene.

In another embodiment, the invention is an isolated nucleic acid molecule comprising a a polynucleotide that hybridizes under stringent conditions to a quorum sensing controlled genetic locus derived from the genome of *Pseudomonas aeruginosa*, wherein the genetic locus comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID

NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36, operatively linked to a reporter gene.

In one embodiment, an isolated nucleic acid molecule of the invention comprises a reporter gene contained in a transposable element.

Accordingly, a further aspect of the invention pertains to a vector comprising an isolated nucleic acid molecule of the invention. In another aspect, the invention provides cells containing an isolated nucleic acid molecule of the invention.

An additional aspect of the invention is a method for identifying a modulator of quorum sensing signaling in bacteria. The method comprises:

providing a cell containing an isolated nucleic acid molecule of the invention, wherein the cell is responsive to a quorum sensing signal molecule such that a detectable signal is generated;

contacting said cell with a quorum sensing signal molecule in the presence and absence of a test compound;

and detecting a change in the detectable signal to thereby identify the test compound as a modulator of quorum sensing signaling in bacteria.

Accordingly, in another aspect, the invention provides a compound identified by a method of the invention which modulates, *e.g.*, inhibits, quorum sensing signaling in *Pseudomonas aeruginosa*. In one embodiment, the compound inhibits quorum sensing signaling in *Pseudomonas aeruginosa* by inhibiting an enzyme involved in the synthesis of a quorum sensing signal molecule, by interfering with quorum sensing signal reception, or by scavenging the quorum sensing signal molecule.

The invention also pertains to a method for identifying quorum sensing controlled genes in a cell, and specifically in one particular human pathogen, *Pseudomonas aeruginosa*. Thus, in one aspect, the invention provides a method for identifying a quorum sensing controlled gene in a cell, the method comprising:

providing a cell which is responsive to a quorum sensing signal molecule such that expression of a quorum sensing controlled gene is modulated, and wherein modulation of the expression of said quorum sensing controlled gene generates a detectable signal;

contacting said cell with a quorum sensing signal molecule;

and detecting a change in the detectable signal to thereby identify a quorum sensing signaling controlled gene.

In one embodiment the cell comprises a reporter gene operatively linked to a quorum sensing controlled gene or a regulatory sequence of a quorum sensing controlled gene, such that modulation of the expression of the quorum sensing controlled gene modulates the transcription of the reporter gene, thereby providing a detectable signal.

- 5 In another embodiment the reporter gene is contained in a transposable element. In yet another embodiment, the quorum sensing signal molecule is produced by a second cell, *e.g.*, a bacterial cell. In a further embodiment, the quorum sensing signal molecule is an autoinducer of said quorum sensing controlled gene, *e.g.*, a homoserine lactone, or an analog thereof.

10

Brief Description of the Drawings

Figure 1 depicts the paradigm for quorum sensing signaling in the target bacterium, *Pseudomonas aeruginosa*.

15

Figure 2 depicts patterns of β -galactosidase expression in representative qsc mutants and in a strain with a *lasB::lacZ* chromosomal fusion generated by site-specific mutation. Units of β -galactosidase are given as a function of culture density for cells grown without added signal molecules (○), with added 3OC₁₂-HSL (●), with added C₄-HSL (■), or with both signals added (□).

20

Figure 3 depicts the nucleic acid sequence of the quorum sensing controlled locus on the *P. aeruginosa* chromosome mapped in the *P. aeruginosa* mutant strain qsc102.

25

Figure 4 depicts putative qsc operons. Open reading frames (ORFs) are indicated by the arrows. ORFs discovered in the qsc screen are indicated by their qsc number.

Figure 5 depicts a growth curve of PAO1/pMW303G. Culture growth is monitored at 600 nm (closed circles) and β -galactosidase activity is measured with a chemiluminescent substrate analog in relative light units (RLU; open circles).

30

Figure 6 is a map of the qsc insertions on the *P. aeruginosa* chromosome. Arrowheads indicate the direction of *lacZ* transcription. In addition to the qsc mutants, *lasR* and *lasI*, *rhlR*, and *lasB* are also mapped. The locations of *las*-boxes like elements are shown as black dots between the two DNA strands. The numbers indicate distance in megabases on the approximately 6 megabase chromosome.

35

Figure 7 depicts putative *las*-type boxes in upstream DNA regions of *qsc* mutants. ORFs as described in Materials and Methods. Bases outlined in black represent residues conserved in all sequences and gray outlines are conserved in 8 of 10 sequences.

5

Figure 8 depicts the principle of a bioassay for modulators of quorum sensing signaling. Strain PAO1 produces the signal 3-oxo-C12-HSL. Strain QSC102 responds by inducing *lacZ*.

10

Figure 9 depicts the results of an assay performed using the test compound acetyl-butyrolactone, which is present in the wells at increasing concentration (mM, as indicated). There are two rows and two columns per concentration to show reproducibility of the assay.

15

Figure 10A depicts the structure of a mobilizable plasmid for generating an indicator strain. Filled boxes represent chromosomal DNA derived from the *P. aeruginosa* locus where *lacZ* is inserted in strain QSC102.

20

Figure 10B depicts induction of β -galactosidase as PAQ1 reaches high density. Cell growth is monitored at 600 nm (closed circles) and expression of β -galactosidase is measured in Miller units (open circles).

Figure 11 depicts the reaction mechanism of the RhlI autoinducer synthase.

25

Figure 12 depicts a continuous culture bioreactor.

Detailed Description of the Invention

In gram-negative bacteria, such as *Pseudomonas aeruginosa*, quorum sensing involves two proteins, the autoinducer synthase - the I protein - and the transcriptional activator - the R protein. The synthase produces an acylated homoserine lactone (the "autoinducer"; see structure **1** below), which can diffuse into the surrounding environment (Fuqua, C. *et al.* (1998) *Curr Opin Microbiol.* 1(2):183-189; Fuqua, *et al.* 1994. *J Bacteriol.* 176(2):269-75). The autoinducer molecule is composed of an acyl chain in a peptide bond with the amino nitrogen of a homoserine lactone (HSL). For different quorum sensing systems, the side-chain may vary in length, degree of saturation, and oxidation state. As the density of bacteria increases, so does the concentration of this freely diffusible signal molecule. Once the concentration reaches a defined threshold, it binds to the R-protein, which then activates transcription of


007060-0225950

numerous genes. Of particular interest are genes involved in pathogenicity and in biofilm formation (see Figure 1).

Pseudomonas aeruginosa has two quorum sensing systems, *las* and *rhl*, named for their role in the expression of elastase, and the RhlI/RhlR proteins, which were first described for their role in rhamnolipid biosynthesis. (Hanzelka, B.A. *et al.* (1996) *J. Bacteriol.* 178:5291-5294; Baldwin, T.O. *et al.* (1989) *J. of Biol. and Chem.* 4:326-341; Passador, L., *et al.* (1993) *Science* 260:1127-1130; Pearson, J.P. *et al.* (1994) *PNAS* 91:197-201; Pesci, E.C. *et al.* (1997) *Trends in Microbiol.* 5(4):132-135; Pesci, E.C. *et al.* (1997) *J. Bacteriol.* 179:3127-3132). The two systems have distinct autoinducer synthases (*lasI* and *rhlI*), transcriptional regulators (*lasR* and *rhlR*), and autoinducers (*N*-(3-oxododecanoyl) homoserine lactone (HSL) and *N*-butyryl HSL) (Sitnikov, D.M. *et al.* (1995) *Mol. Microbiol.* 17:801-812). The *rhl* and *las* systems are involved in regulating the expression of a number of secreted virulence factors, biofilm development, and the stationary phase sigma factor (RpoS) (Davies, D.G. *et al.* (1998) *Science* 280:295-298; Latifi, A. *et al.* (1995) *Mol. Microbiol. Rev.* 17:333-344; Ochsner, U.A., *et al.* (1995) *PNAS*, 92:6424-6428; Pesci, E.C. *et al.* (1997) *Trends in Microbiol.* 5(4):132-135; Pesci, E.C. *et al.* (1997) *J. Bacteriol.* 179:3127-3132). Expression of the *rhl* system requires a functional *las* system, therefore the two systems in combination with RpoS constitute a regulatory cascade (Pesci, E.C. *et al.* (1997) *Trends in Microbiol.* 5(4):132-135; Pesci, E.C. *et al.* (1997) *J. Bacteriol.* 179:3127-3132, Seed *et al.* 1995).

The signal in the Las system is 3-oxo-dodecanoyl-HSL (3-oxo-C12-HSL) **2**, while the signal used in the Rhl system is butanoyl-HSL (C4-HSL) **3**. It has been shown that 3-oxo-C12-HSL increases expression of RhlR, indicating a hierarchy of regulation systems (Pesci, E. C. *et al.* (1997) *Trends Microbiol.* 5(4):132-4). The Las signal 3-oxo-C12-HSL is synthesized by LasI along with a small amount of *N*-(3-oxooctanoyl) HSL and *N*-(3-oxohexanoyl) HSL, while RhlI makes primarily the signal C4-HSL and a small amount of *N*-hexanoyl (Pearson, J.P. *et al.* (1997) *J. Bacteriol.* 179:5756-5757; Winson, M.K. *et al.* (1995) *PNAS* 92:9427-9431).

30



Bacterial signaling triggers the expression of a number of virulence factors in *P. aeruginosa* including two elastases, an alkaline protease and exotoxin A (Pesci, E. C. *et al.* (1997) *Trends Microbiol.* 5(4):132-4; Pesci, E. C. *et al.* (1997) *J Bacteriol.*

179(10):3127-32) - proteins that allow the organism to attack host tissue. Bacterial signaling also controls the expression of the antioxidant pyocyanin, a compound that allows the bacteria to neutralize one important host defense, the generation of superoxide radicals (Britigan, *et al.* (1999) *Infect Immun.* 67(3):1207-12, Hassan, H. M. *et al.* (1979) *Arch Biochem Biophys.* 196(2):385-95, Hassan, H. M. *et al.* 1980. *J Bacteriol.* 141(1):156-63). It has been shown in a neonatal mouse model that a defined mutant of *P. aeruginosa* which lacks the signal receptor protein (LasR) was significantly less virulent than the wild type PAO1, as measured by the ability to cause acute pneumonia, bacteremia and death (Tang, H. B. *et al.* (1996) *Infect Immun.* 64(1):37-43).

The invention is based on the interruption of bacterial cell-to-cell signaling, *i.e.*, quorum sensing signaling in order to render a bacterial population more susceptible to treatment, either through the host immune-response or in combination with traditional antibacterial agents and biocides. Thus, the invention provides a bacterial indicator strain that allows for a high throughput screening assay for identifying compounds that modulate, *e.g.*, inhibit bacterial cell-to-cell signalling. The compounds so identified will provide novel anti-pathogenics and anti-fouling agents.

Definitions

Before further description of the invention, certain terms employed in the specification, examples and appended claims are, for convenience, collected here.

The term "analog" as in "homoserine lactone analog" is intended to encompass compounds that are chemically and/or electronically similar but have different atoms, such as isosteres and isologs. An analog includes a compound with a structure similar to that of another compound but differing from it in respect to certain components or structural makeup. The term analog is also intended to encompass stereoisomers.

The language "autoinducer compounds" is art-recognized and is intended to include molecules, *e.g.*, proteins which freely diffuse across cell membranes and which activate transcription of various factors which affect bacterial viability. Such compounds can affect virulence, and biofilm development. Autoinducer compounds can be acylated homoserine lactones. They can be other compounds similar to those listed in Table 1. Homoserine autoinducer compounds are produced *in vivo* by the interaction of a homoserine lactone substrate and an acylated acyl carrier protein in a reaction catalyzed by an autoinducer synthase molecule. In isolated form, autoinducer compounds can be obtained from naturally occurring proteins by purifying cellular extracts, or they can be chemically synthesized or recombinantly produced.

The language "autoinducer synthase molecule" is intended to include molecules, *e.g.* proteins, which catalyze or facilitate the synthesis of autoinducer compounds, *e.g.* in the quorum sensing system of bacteria. It is also intended to include active portions of the

[illegible]

Table 1 contains a list of exemplary autoinducer synthase proteins of the quorum sensing systems of various gram-negative bacteria.

5

Table 1. Summary of *N*-acyl homoserine lactone based regulatory systems

Bacterial species	Signal molecules ^a	Regulatory Proteins ^b	Target function(s)
<i>Vibrio fischeri</i>	N-3-(oxohexanoyl)-homoserine lactone (VAI-1)	LuxI/LuxR	<i>lux</i> /CDABEG, <i>luxR</i> luminescence
	N-(octanoyl)-L-homoserine lactone (VAI-2)	AinS/AinR ^c	<i>lux</i> /CDABEG, ?
<i>Vibrio harveyi</i>	<i>N</i> - β -(hydroxybutyryl)-homoserine lactone (HAI-1)	LuxM/LuxN-LuxO-LuxR ^d	<i>lux</i> /CDABEG, luminescence and polyhydroxybutyrate synthesis
	HAI-2	Lux [?] /LuxPQ-LuxO-LuxR ^d	<i>lux</i> CDABEG
<i>Pseudomonas aeruginosa</i>	N-3-(oxododecanoyl)-L-homoserine lactone (PAI-1)	LasI/LasR	<i>lasB</i> , <i>lasA</i> , <i>aprA</i> , <i>toxA</i> , virulence factors
	N-(butyryl)-L-homoserine lactone (PAI-2)	RhlI/RhlR	<i>rhlAB</i> , rhamnolipid synthesis, virulence factors
<i>Pseudomonas aeureofaciens</i>	(PRAI) ^e	PhzI/PhzR	<i>phz</i> , phenazine biosynthesis
<i>Agroacterium tumefaciens</i>	N-3-(oxooctanoyl)-L-homoserine lactone (AAI)	TraI/TraR-TraM	<i>tra</i> gens, <i>traR</i> , Ti plasmid conjugal transfer
<i>Erwinia carotovora</i> subsp. <i>carotovora</i> SCRI193	VAI-1 ^f	ExpI/ExpR	<i>pel</i> , <i>pec</i> , <i>pep</i> , exoenzyme synthesis
<i>Erwinia carotovora</i> subsp. <i>carotovora</i> SCC3193	VAI-1 ^f	CarI/CarR	<i>cap</i> , carbapenem antibiotic synthesis
<i>Erwinia carotovora</i> subsp. <i>carotovora</i> 71	VAI-1 ^f	HsII/?	<i>pel</i> , <i>pec</i> , <i>pep</i> , exoenzyme synthesis
<i>Erwinia stewartii</i>	VAI-1 ^f	EsaI/EsaR	<i>wts</i> genes, exopolysaccharide synthesis, virulence factors
<i>Rhizobium leguminosarum</i>	<i>N</i> -(3R-hydroxy-7- <i>cis</i> -tetradecanoyl)-L-homoserine lactone, small bacteriocin, (RLAI)	?/RhlR	<i>rhlABC</i> , rhizosphere genes and stationary phase
<i>Enterobacter agglomerans</i>	VAI-1 ^f	EagI/EagR	function unclear
<i>Yersenia enterocolitica</i>	VAI-1 ^f	YenI/YenR	function unclear
<i>Serratia liquifaciens</i>	<i>N</i> -butanoyl-L-homoserine lacton (SAI-1)	SwrI/?	swarming motility
	<i>N</i> -hexanoyl-L-homoserine lacton (SAI-2)	SwrI/?	swarming motility
<i>Aeromonas hydrophila</i>	(AHAI) ^e	AhyI/AhyR	function unclear
<i>Escherichia coli</i> /?? ^g		?/SdiA	<i>ftsQAZ</i> , cell division

Autoinducer synthase molecules can be obtained from naturally occurring sources, *e.g.*, by purifying cellular extracts, can be chemically synthesized or can be recombinantly produced. Recombinantly produced autoinducer synthase molecules can have the amino acid sequence of a naturally occurring form of the autoinducer synthase protein. They can also have a similar amino acid sequence which includes mutations such as substitutions and deletions (including truncation) of a naturally occurring form of the protein. Autoinducer synthase molecules can also include molecules which are structurally similar to the structures of naturally occurring autoinducer synthase proteins, *e.g.*, biologically active variants.

TraI, LuxI, RhII are the homoserine lactone autoinducer synthases of *Agrobacterium tumefaciens*, *Vibrio fischeri*, and *Pseudomonas aeruginosa*, respectively. The term "RhII" is intended to include proteins which catalyze the synthesis of the homoserine lactone autoinducer of the RhII quorum sensing system of *P. aeruginosa*, butyryl homoserine lactone.

The term "biofilm" is intended to include biological films that develop and persist at interfaces in aqueous environments. Biofilms are composed of microorganisms embedded in an organic gelatinous structure composed of one or more matrix polymers which are secreted by the resident microorganisms. The language "biofilm development" or "biofilm formation" is intended to include the formation, growth, and modification of the bacterial colonies contained within the biofilm structures as well as the synthesis and maintenance of the exopolysaccharide matrix of the biofilm structures.

The term "compound" as used herein (*e.g.*, as in "test compound," or "modulator compound") is intended to include both exogenously added test compounds and peptides endogenously expressed from a peptide library. Test compounds may be purchased, chemically synthesized or recombinantly produced. Test compounds can be obtained from a library of diverse compounds based on a desired activity, or alternatively they can be selected from a random screening procedure. In one embodiment, an indicator cell (*e.g.*, a cell which responds to quorum sensing signals by generating a detectable signal) also produces the test compound which is being screened. For instance, the indicator cell can produce, *e.g.*, a test polypeptide, a test nucleic acid and/or a test carbohydrate, which is screened for its ability to modulate quorum sensing signaling. In such embodiments, a culture of such reagent cells will collectively provide a library of potential modulator molecules and those members of the library which either stimulate or inhibit quorum sensing signaling can be selected and identified. In another embodiment, a test compound is produced by a second cell which is co-incubated with the indicator cell.

The terms “derived from” or “derivative”, as used interchangeably herein, are intended to mean that a sequence is identical to or modified from another sequence, *e.g.*, a naturally occurring sequence. Derivatives within the scope of the invention include polynucleotide derivatives. Polynucleotide or nucleic acid derivatives differ from the sequences described herein (*e.g.*, SEQ ID Nos.: 1-38) or known in nucleotide sequence. For example, a polynucleotide derivative may be characterized by one or more nucleotide substitutions, insertions, or deletions, as compared to a reference sequence. A nucleotide sequence comprising a quorum sensing controlled genetic locus that is derived from the genome of *P. aeruginosa*, *e.g.*, SEQ ID Nos.: 1-38, includes sequences that have been modified by various changes such as insertions, deletions and substitutions, and which retain the property of being regulated in response to a quorum sensing signaling event. Such sequences may comprise a quorum sensing controlled regulatory element and/or a quorum sensing controlled gene. The nucleotide sequence of the *P. aeruginosa* genome is available at www.pseudomonas.com.

Polypeptide or protein derivatives include polypeptide or protein sequences that differ from the sequences described or known in amino acid sequence, or in ways that do not involve sequence, or both, and still preserve the activity of the polypeptide or protein. Derivatives in amino acid sequence are produced when one or more amino acids is substituted with a different natural amino acid, an amino acid derivative or non-native amino acid. In certain embodiments protein derivatives include naturally occurring polypeptides or proteins, or biologically active fragments thereof, whose sequences differ from the wild type sequence by one or more conservative amino acid substitutions, which typically have minimal influence on the secondary structure and hydrophobic nature of the protein or peptide. Derivatives may also have sequences which differ by one or more non-conservative amino acid substitutions, deletions or insertions which do not abolish the biological activity of the polypeptide or protein.

Conservative substitutions (substituents) typically include the substitution of one amino acid for another with similar characteristics (*e.g.*, charge, size, shape, and other biological properties) such as substitutions within the following groups: valine, glycine; glycine, alanine; valine, isoleucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. The non-polar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

In other embodiments, derivatives with amino acid substitutions which are less conservative may also result in desired derivatives, *e.g.*, by causing changes in charge,

U.S. Pat. No. 7,000,000

conformation and other biological properties. Such substitutions would include, for example, substitution of hydrophilic residue for a hydrophobic residue, substitution of a cysteine or proline for another residue, substitution of a residue having a small side chain for a residue having a bulky side chain or substitution of a residue having a net positive charge for a residue having a net negative charge. When the result of a given substitution cannot be predicted with certainty, the derivatives may be readily assayed according to the methods disclosed herein to determine the presence or absence of the desired characteristics. The polypeptides and proteins of this invention may also be modified by various changes such as insertions, deletions and substitutions, either conservative or nonconservative where such changes might provide for certain advantages in their use.

As used herein, the term "genetic locus" includes a position on a chromosome, or within a genome, which is associated with a particular gene or genetic sequences having a particular characteristic. For example, in one embodiment, a quorum sensing controlled genetic locus includes nucleic acid sequences which comprise an open reading frame (ORF) of a quorum sensing controlled gene. In another embodiment, a quorum sensing controlled genetic locus includes nucleic acid sequences which comprise transcriptional regulatory sequences that are responsive to quorum sensing signaling (*e.g.*, a quorum sensing controlled regulatory element). Examples of quorum sensing controlled genetic loci of *P. aeruginosa* are described herein as SEQ ID NOs.:1-38.

The term "modulator", as in "modulator of quorum sensing signaling" is intended to encompass, in its various grammatical forms, induction and/or potentiation, as well as inhibition and/or downregulation of quorum sensing signaling and/or quorum sensing controlled gene expression. As used herein, the term "modulator of quorum sensing signaling" includes a compound or agent that is capable of modulating or regulating at least one quorum sensing controlled gene or quorum sensing controlled genetic locus, *e.g.*, a quorum sensing controlled genetic locus in *P. aeruginosa*, as described herein. A modulator of quorum sensing signaling may act to modulate either signal generation (*e.g.*, the synthesis of a quorum sensing signal molecule), signal reception (*e.g.*, the binding of a signal molecule to a receptor or target molecule), or signal transmission (*e.g.*, signal transduction via effector molecules to generate an appropriate biological response). In one embodiment, a method of the present invention encompasses the modulation of the transcription of an indicator gene in response to an autoinducer molecule. In another embodiment, a method of the present invention encompasses the modulation of the transcription of an indicator gene, preferably an quorum sensing controlled indicator gene, by a test compound.

The term “operatively linked” or “operably linked” is intended to mean that molecules are functionally coupled to each other in that the change of activity or state of one molecule is affected by the activity or state of the other molecule. In one embodiment, nucleotide sequences are "operatively linked" when the regulatory sequence functionally relates to the DNA sequence encoding the polypeptide or protein of interest. For example, a nucleotide sequence comprising a transcriptional regulatory element(s) (*e.g.*, a promoter) is operably linked to a DNA sequence encoding the protein or polypeptide of interest if the promoter nucleotide sequence controls the transcription of the DNA sequence encoding the protein of interest. In addition, two nucleotide sequences are operatively linked if they are coordinately regulated and/or transcribed. Typically, two polypeptides that are operatively linked are covalently attached through peptide bonds.

The term “quorum sensing signaling” or “quorum sensing” is intended to include the generation of a cellular signal in response to cell density. In one embodiment, quorum sensing signaling mediates the coordinated expression of specific genes. A “quorum sensing controlled gene” is any gene, the expression of which is regulated in a cell density dependent fashion. In a preferred embodiment, the expression of a quorum sensing controlled gene is modulated by a quorum sensing signal molecule, *e.g.*, an autoinducer molecule (*e.g.*, a homoserine lactone molecule). The term “quorum sensing signal molecule” is intended to include a molecule that transduces a quorum sensing signal and mediates the cellular response to cell density. In a preferred embodiment the quorum sensing signal molecule is a freely diffusible autoinducer molecule, *e.g.*, a homoserine lactone molecule or analog thereof. In one embodiment, a quorum sensing controlled gene encodes a virulence factor. In another embodiment, a quorum sensing controlled gene encodes a protein or polypeptide that, either directly or indirectly, inhibits and/or antagonizes a bacterial host defense mechanism. In yet another embodiment, a quorum sensing controlled gene encodes a protein or polypeptide that regulates biofilm formation.

The term “regulatory sequences” is intended to include the DNA sequences that control the transcription of an adjacent gene. Gene regulatory sequences include, but are not limited to, promoter sequences that are found in the 5' region of a gene proximal to the transcription start site which bind RNA polymerase to initiate transcription. Gene regulatory sequences also include enhancer sequences which can function in either orientation and in any location with respect to a promoter, to modulate the utilization of a promoter, and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel (1990) *Methods Enzymol.* 185:3-7. Transcriptional control elements include, but are not limited to, promoters, enhancers, and repressor and activator binding sites. The gene regulatory sequences of

the present invention contain binding sites for transcriptional regulatory proteins. In one embodiment, a regulatory sequence includes a sequence that mediates quorum sensing controlled gene expression, *e.g.*, a las box. In a preferred embodiment, gene regulatory sequences comprise sequences derived from the *Pseudomonas aeruginosa* genome which modulate quorum sensing controlled gene expression, *e.g.*, SEQ ID NOs.:38 and 39. In another preferred embodiment, gene regulatory sequences comprise sequences (*e.g.*, a genetic locus) derived from the *Pseudomonas aeruginosa* genome which modulate the expression of quorum sensing controlled genes, *e.g.*, SEQ ID NOs.:1-36.

The term "reporter gene" or "indicator gene" generically refers to an expressible (*e.g.*, able to be transcribed and (optionally) translated) DNA sequence which is expressed in response to the activity of a transcriptional regulatory protein. Indicator genes include unmodified endogenous genes of the host cell, modified endogenous genes, or a reporter gene of a heterologous construct, *e.g.*, as part of a reporter gene construct. In a preferred embodiment, the level of expression of an indicator gene produces a detectable signal.

Reporter gene constructs are prepared by operatively linking an indicator gene with at least one transcriptional regulatory element. If only one transcriptional regulatory element is included, it is advantageously a regulatable promoter. In a preferred embodiment at least one of the selected transcriptional regulatory elements is directly or indirectly regulated by quorum sensing signals, whereby quorum sensing controlled gene expression can be monitored via transcription and/or translation of the reporter genes.

Many reporter genes and transcriptional regulatory elements are known to those of skill in the art and others may be identified or synthesized by methods known to those of skill in the art. Reporter genes include any gene that expresses a detectable gene product, which may be RNA or protein. Preferred reporter genes are those that are readily detectable. In one embodiment, an indicator gene of the present invention is comprised in the nucleic acid molecule in the form of a fusion gene (*e.g.*, operatively linked) with a nucleotide sequence that includes regulatory sequences (*e.g.*, quorum sensing transcriptional regulatory elements, *e.g.*, a las box) derived from the *Pseudomonas aeruginosa* genome (*e.g.*, SEQ ID NOs.:38 and 39). In another embodiment, an indicator gene of the present invention is operatively linked to quorum sensing transcriptional regulatory sequences that regulate a quorum sensing controlled genetic locus derived from the *Pseudomonas aeruginosa* genome, *e.g.*, a genetic locus comprising a nucleotide sequence set forth as SEQ ID NOs.: 1-36. In yet another embodiment, an indicator gene of the present invention is operatively linked to a nucleotide sequence comprising a quorum sensing controlled genetic locus derived from the *Pseudomonas aeruginosa* genome (*e.g.*, SEQ ID NOs.:1-39). In certain

embodiments of the invention, an indicator gene (*e.g.*, a promoterless indicator gene) is contained in a transposable element.

The term "detecting a change in the detectable signal" is intended to include the detection of alterations in gene transcription of an indicator or reporter gene induced upon modulation of quorum sensing signaling. In certain embodiments, the reporter gene may provide a selection method such that cells in which the transcriptional regulatory protein activates transcription have a growth advantage. For example the reporter could enhance cell viability, relieve a cell nutritional requirement, and/or provide resistance to a drug. In other embodiments, the detection of an alteration in a signal produced by an indicator gene encompass assaying general, global changes to the cell such as changes in second messenger generation.

The amount of transcription from the reporter gene may be measured using any method known to those of skill in the art. For example, specific mRNA expression may be detected using Northern blots, or a specific protein product may be identified by a characteristic stain or an intrinsic activity. In preferred embodiments, the gene product of the reporter is detected by an intrinsic activity associated with that product. For instance, the reporter gene may encode a gene product that, by enzymatic activity, gives rise to a detection signal based on color, fluorescence, or luminescence.

The amount of regulation of the indicator gene, *e.g.*, expression of a reporter gene, is then compared to the amount of expression in a control cell. For example, the amount of transcription of an indicator gene may be compared between a cell in the absence of a test modulator molecule and an identical cell in the presence of a test modulator molecule.

As used interchangeably herein, the terms "transposon" and "transposable element" are intended to include a piece of DNA that can insert into and cut itself out of, genomic DNA of a particular host species. Transposons include mobile genetic elements (MGEs) containing insertion sequences and additional genetic sequences unrelated to insertion functions (for example, sequences encoding a reporter gene). Insertion sequence elements include sequences that are between 0.7 and 1.8 kb in size with termini approximately 10 to 40 base pairs in length with perfect or nearly perfect repeats. As used herein, a transposable element is operatively linked to the nucleotide sequence into which it is inserted. Transposable elements are well known in the art, and are described for example, at www.bact.wisc.edu/MicrotextBook/BactGenetics.

The present invention discloses a method for identifying modulators of quorum sensing signaling in bacteria, *e.g.*, *Pseudomonas aeruginosa*. As described herein, the method of the invention comprises providing a cell which comprises a quorum sensing controlled gene, wherein the cell is responsive to a quorum sensing signal molecule such that a detectable signal is generated. A cell which responds to a quorum sensing signal

5 molecule by generating a detectable signal is referred to herein as an “indicator cell” or a “reporter cell”. In a preferred embodiment of the invention, the cell is a *P. aeruginosa* bacterial cell. In another preferred embodiment, the cell is from a mutant strain of *P. aeruginosa* which comprises a reporter gene operatively linked to a regulatory sequence of a quorum sensing controlled gene, wherein said mutant strain is responsive to a quorum sensing signal molecule, such that a detectable signal is generated. In yet another preferred embodiment, the cell is a mutant strain of *P. aeruginosa* which comprises a promoterless reporter gene inserted in the chromosome at a quorum sensing controlled genetic locus, *e.g.*, a genetic locus comprising a nucleotide sequence set forth as SEQ ID NOs.:1-38, wherein said mutant strain is responsive to a quorum sensing signal molecule such that a detectable signal is generated by the reporter gene. In a preferred embodiment, the reporter gene is contained in a transposable element. In a further preferred embodiment, the cell is from a strain of *P. aeruginosa* in which *lasI* and *rhlI* are inactivated, such that the cell does not express the *lasI* and *rhlI* autoinducer synthases which are involved in the generation of quorum sensing signal molecules. A compound is identified as a modulator of quorum sensing signaling in bacteria by contacting the cell with a quorum sensing signal molecule in the presence and absence of a test compound and detecting a change in the detectable signal.

20 Quorum sensing signal molecules that are useful in the methods of the present invention include autoinducer compounds such as homoserine lactones, and analogs thereof (see Table 1). In certain embodiments, the quorum sensing signal molecule is either 3-oxo-C12-homoserine lactone or C4-HSL. In one embodiment, the cell does not express the quorum sensing signal molecule. For example, the cell may comprise a mutant strain of *Pseudomonas aeruginosa* wherein *lasI* and *rhlI* are inactivated. Therefore, the cell is contacted with an exogenous quorum sensing signal molecule, *e.g.*, a recombinant or synthetic molecule. In another embodiment, the quorum sensing signal molecule is produced by a second cell (*e.g.*, a prokaryotic or eukaryotic cell), which is co-incubated with the indicator cell. For example, an indicator cell which does not express a quorum sensing signal molecule can be co-incubated with a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule. Alternatively, the indicator strain which does not express a quorum sensing signal molecule is co-incubated with a second cell which has been transformed, or otherwise altered, such that it is able to express a quorum sensing signal molecule. In yet another embodiment, the quorum sensing signal molecule is expressed by the indicator strain.

35 Similarly, the test compound can be exogenously added to an indicator strain, produced by a second cell which is co-incubated with the indicator strain, or expressed by the indicator strain. Exemplary compounds which can be screened for activity

include, but are not limited to, peptides, nucleic acids, carbohydrates, small organic molecules, and natural product extract libraries.

The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including:

5 biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries

10 of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:45).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example, in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

15

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

20

In certain embodiments of the instant invention, the compounds tested are in the form of peptides from a peptide library. The peptide library may take the form of a cell culture, in which essentially each cell expresses one, and usually only one, peptide of the library. While the diversity of the library is maximized if each cell produces a peptide of a different sequence, it is usually prudent to construct the library so there is some redundancy. Depending on size, the combinatorial peptides of the library can be

25 expressed as is, or can be incorporated into larger fusion proteins. The fusion protein can provide, for example, stability against degradation or denaturation. In an exemplary embodiment of a library for intracellular expression, *e.g.*, for use in conjunction with intracellular target receptors, the polypeptide library is expressed as thioredoxin fusion proteins (see, for example, U.S. Patents 5,270,181 and 5,292,646; and PCT publication WO94/ 02502). The combinatorial peptide can be attached on the terminus of the

30 thioredoxin protein, or, for short peptide libraries, inserted into the so-called active loop.

35

In one embodiment of the instant invention the cell further comprises a means for generating the detectable signal. For example, the cell may comprise a reporter

gene, the transcription of which is regulated by a quorum sensing signal molecule. In a preferred embodiment, the reporter gene is operatively linked to a regulatory sequence of a quorum sensing controlled gene, *e.g.* a nucleotide sequence comprising at least one quorum sensing controlled regulatory element, *e.g.*, a *las* box. In another embodiment, the reporter gene is operatively linked to a quorum sensing controlled genetic locus, *e.g.*, a quorum sensing controlled gene, such that transcription of the indicator gene is responsive to quorum sensing signals. For example, in a preferred embodiment, a promoterless reporter gene is inserted into a quorum sensing controlled genetic locus derived from the genome of *P. aeruginosa*. Such quorum sensing controlled genetic loci, as described herein, include the loci in the *P. aeruginosa* genome which comprise the nucleotide sequences set forth as SEQ ID NOs.: 1-38. In another preferred embodiment, the promoterless reporter gene is contained in a transposable element that is inserted into a quorum sensing controlled genetic locus in the *P. aeruginosa* genome.

Examples of reporter genes include, but are not limited to, CAT (chloramphenicol acetyl transferase) (Alton and Vapnek (1979), *Nature* 282: 864-869), and other enzyme detection systems, such as beta-galactosidase (*lacZ*), firefly luciferase (deWet *et al.* (1987), *Mol. Cell. Biol.* 7:725-737); bacterial luciferase (Engebrecht and Silverman (1984), *PNAS* 1: 4154-4158; Baldwin *et al.* (1984), *Biochemistry* 23: 3663-3667); alkaline phosphatase (Toh *et al.* (1989) *Eur. J. Biochem.* 182: 231-238, Hall *et al.* (1983) *J. Mol. Appl. Gen.* 2: 101), human placental secreted alkaline phosphatase (Cullen and Malim (1992) *Methods in Enzymol.* 216:362-368), and horseradish peroxidase. In one preferred embodiment, the indicator gene is *lacZ*. In another preferred embodiment, the indicator gene is green fluorescent protein (U.S. patent 5,491,084; WO96/23898) or a variant thereof. A preferred variant is GFPmut2. Other reporter genes include *ADE1*, *ADE2*, *ADE3*, *ADE4*, *ADE5*, *ADE7*, *ADE8*, *ASP3*, *ARG1*, *ARG3*, *ARG4*, *ARG5*, *ARG6*, *ARG8*, *ARO2*, *ARO7*, *BAR1*, *CAT*, *CHO1*, *CYS3*, *GAL1*, *GAL7*, *GAL10*, *HIS1*, *HIS3*, *HIS4*, *HIS5*, *HOM3*, *HOM6*, *ILV1*, *ILV2*, *ILV5*, *INO1*, *INO2*, *INO4*, *LEU1*, *LEU2*, *LEU4*, *LYS2*, *MAL*, *MEL*, *MET2*, *MET3*, *MET4*, *MET8*, *MET9*, *MET14*, *MET16*, *MET19*, *OLE1*, *PHO5*, *PRO1*, *PRO3*, *THR1*, *THR4*, *TRP1*, *TRP2*, *TRP3*, *TRP4*, *TRP5*, *URA1*, *URA2*, *URA3*, *URA4*, *URA5* and *URA10*.

In accordance with the methods of the invention, compounds which modulate quorum sensing signaling can be selected and identified. The ability of compounds to modulate quorum sensing signaling can be detected by up or down-regulation of the detection signal provided by the indicator gene. Any difference, *e.g.*, a statistically significant difference, in the amount of transcription indicates that the test compound has in some manner altered the activity of quorum sensing signaling.

A modulator of quorum sensing signaling may act by inhibiting an enzyme involved in the synthesis of a quorum sensing signal molecule, by inhibiting reception of

the quorum sensing signal molecule by the cell, or by scavenging the quorum sensing signal molecule. The term “scavenging” is meant to include the sequestration, chemical modification, or inactivation of a quorum sensing signal molecule such that it is no longer able to regulate quorum sensing gene control. After identifying certain test compounds as potential modulators of quorum sensing signaling, the practitioner of the subject assay will continue to test the efficacy and specificity of the selected compounds both *in vitro* and *in vivo*, *e.g.*, in an assay for bacterial viability and/or pathogenicity.

In another aspect, the present invention discloses a method for identifying a quorum sensing controlled gene in bacteria, *e.g.*, *Pseudomonas aeruginosa*. The method comprises providing a cell which is responsive to a quorum sensing signal molecule such that expression of a quorum sensing controlled gene is modulated, and wherein modulation of the expression of the quorum sensing controlled gene generates a detectable signal. The cell is contacted with a quorum sensing signal molecule and a change in the signal is detected to thereby identify a quorum sensing signaling controlled gene.

In one embodiment, the cell further comprises a means for generating the detectable signal, *e.g.*, a reporter gene. For example, the cell may comprise a promoterless reporter gene that is operatively linked to a quorum sensing controlled genetic locus such that modulation of the expression of the quorum sensing controlled locus concurrently modulates transcription of the reporter gene. The position of the quorum sensing controlled genetic locus is then mapped based on the position of the reporter gene.

In a preferred embodiment of the invention, the cell is a *P. aeruginosa* bacterial cell. In another preferred embodiment, the cell is a mutant strain of *P. aeruginosa* which comprises a promoterless reporter gene inserted in the chromosome at a quorum sensing controlled genetic locus, *e.g.*, a genetic locus comprising a nucleotide sequence set forth as SEQ ID NOs.:1-39, wherein said mutant strain is responsive to a quorum sensing signal molecule such that a detectable signal is generated by the reporter gene. In a preferred embodiment, the reporter gene is contained in a transposable element. In a further preferred embodiment, the cell is from a strain of *P. aeruginosa* in which *lasI* and *rhlI* are inactivated, such that the cell does not express the *lasI* and *rhlI* autoinducer synthases which are involved in the generation of quorum sensing signal molecules.

It is also to be understood that genomic sequences from a mutant bacterial strain (*e.g.*, *P. aeruginosa*) in which a promoterless reporter gene (*e.g.*, a reporter gene contained in a transposable element) has been inserted at a quorum sensing controlled locus, can be assayed in a heterologous cell that is responsive to a quorum sensing signal molecule such that quorum sensing signal transduction occurs. For example, the genomic DNA of a strain of *P. aeruginosa* subjected to transposon mutagenesis, as

007060:0225950

described herein, can be engineered into a library, and transferred to another cell capable of quorum sensing signaling (e.g., a different species of gram negative bacteria), and assayed to identify a quorum sensing controlled gene.

In one embodiment, the cell is contacted with an exogenous quorum sensing signal molecule, e.g., a recombinant or synthetic molecule, as described herein. In another embodiment, the quorum sensing signal molecule is produced by a second cell (e.g., a prokaryotic or eukaryotic cell), which is co-incubated with the indicator cell. For example, an indicator cell which does not express a quorum sensing signal molecule can be co-incubated with a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule. Alternatively, the indicator strain which does not express a quorum sensing signal molecule is co-incubated with a second cell which has been transformed, or otherwise altered, such that it is able to express a quorum sensing signal molecule. In yet another embodiment, the quorum sensing signal molecule is expressed by the indicator strain.

Another aspect of the invention provides a mutant strain of *Pseudomonas aeruginosa* comprising a promoterless reporter gene inserted in a chromosome at a genetic locus comprising a nucleotide sequence set forth as SEQ ID NOs:1-36, e.g., a quorum sensing controlled genetic locus. In one embodiment the reporter gene is contained in a transposable element. In another embodiment, the reporter gene is lacZ or GFP, or a variant thereof, e.g., GFPmut2. In yet another embodiment, *lasI* and *rhlI* are inactivated in the mutant strain of *P. aeruginosa*. The above-described cells are useful in the methods of the instant invention, as the cells are responsive to a quorum sensing signal molecule such that a detectable signal is generated by the reporter gene. These cells are also useful for studying the function of polypeptides encoded by the quorum sensing controlled loci comprising the nucleotide sequences set forth as SEQ ID NOs.:1-36.

Yet another aspect of the invention provides isolated nucleic acid molecules comprising a nucleotide sequence comprising a quorum sensing controlled genetic locus derived from the genome of *Pseudomonas aeruginosa* operatively linked to a reporter gene. In one embodiment, a reporter gene is operatively linked to a regulatory sequence derived from the genome of *P. aeruginosa*, wherein the regulatory sequence regulates a quorum sensing controlled genetic locus comprising a nucleotide sequence set forth as SEQ ID NO:1-36. In a preferred embodiment such regulatory sequences comprise at least one binding site for a quorum sensing controlled transcriptional regulatory factor (e.g., a transcriptional activator or repressor molecule) such that transcription of the reporter gene is responsive to a quorum sensing signal molecule and/or a modulator of quorum sensing signaling. In another embodiment, a reporter gene is operatively linked to a quorum sensing controlled genetic locus derived from the genome of *P. aeruginosa*,

other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% identical to each other remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, Ausubel *et al.*, eds., John Wiley & Sons, Inc. (1995), sections 2, 4, and 6. Additional stringent conditions can be found in *Molecular Cloning: A Laboratory Manual*, Sambrook *et al.*, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), chapters 7, 9, and 11. A preferred, non-limiting example of stringent hybridization conditions includes hybridization in 4X sodium chloride/sodium citrate (SSC), at about 65-70°C (or alternatively hybridization in 4X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 1X SSC, at about 65-70°C. A preferred, non-limiting example of highly stringent hybridization conditions includes hybridization in 1X SSC, at about 65-70°C (or alternatively hybridization in 1X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 0.3X SSC, at about 65-70°C. A preferred, non-limiting example of reduced stringency hybridization conditions includes hybridization in 4X SSC, at about 50-60°C (or alternatively hybridization in 6X SSC plus 50% formamide at about 40-45°C) followed by one or more washes in 2X SSC, at about 50-60°C. Ranges intermediate to the above-recited values, *e.g.*, at 65-70°C or at 42-50°C are also intended to be encompassed by the present invention. SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1X SSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes each after hybridization is complete. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}\text{C}) = 2(\# \text{ of A} + \text{T bases}) + 4(\# \text{ of G} + \text{C bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G} + \text{C}) - (600/\text{N})$, where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer ($[\text{Na}^+]$ for 1X SSC = 0.165 M). It will also be recognized by the skilled practitioner that additional reagents may be added to hybridization and/or wash buffers to decrease non-specific hybridization of nucleic acid molecules to membranes, for example, nitrocellulose or nylon membranes, including but not limited to blocking agents (*e.g.*, BSA or salmon or herring sperm carrier DNA), detergents (*e.g.*, SDS), chelating agents (*e.g.*, EDTA), Ficoll, PVP and the like. When using nylon membranes, in particular, an additional preferred, non-limiting example of stringent hybridization conditions is hybridization in 0.25-0.5M NaH₂PO₄, 7% SDS at about 65°C, followed by one or more washes at 0.02M NaH₂PO₄, 1% SDS at 65°C (see

e.g., Church and Gilbert (1984) *Proc. Natl. Acad. Sci. USA* 81:1991-1995), or alternatively 0.2X SSC, 1% SDS.

The invention further encompasses nucleic acid molecules that differ from the quorum sensing controlled genetic loci described herein, *e.g.*, the nucleotide sequences shown in SEQ ID NO:1-36. Accordingly, the invention also includes variants, *e.g.*, allelic variants, of the disclosed polynucleotides or proteins; that is naturally occurring and non-naturally occurring alternative forms of the isolated polynucleotide which may also encode proteins which are identical, homologous or related to that encoded by the polynucleotides of the invention.

Nucleic acid variants can be naturally occurring, such as allelic variants (same locus), homologues (different locus), and orthologues (different organism) or can be non naturally occurring. Non-naturally occurring variants can be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. The variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions (as compared in the encoded product). Allelic variants result, for example, from DNA sequence polymorphisms within a population (*e.g.*, a bacterial population) that lead to changes in the nucleic acid sequences of quorum sensing controlled genetic loci.

To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, 90% or 95% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined

5 another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the
10 algorithm of E. Meyers and W. Miller (*Comput. Appl. Biosci.*, 4:11-17 (1988) which has been incorporated into the ALIGN program (version 2.0) (available at <http://vega.igh.cnrs.fr/bin/align-guess.cgi>), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Additionally, the "Clustal" method (Higgins and Sharp, *Gene*, 73:237-44, 1988) and "Megalign" program (Clewley and Arnold, *Methods Mol. Biol.*, 70:119-29, 1997) can be used to align sequences and determine similarity, identity, or homology.

30 Accordingly, the present invention also discloses recombinant vector constructs and recombinant host cells transformed with said constructs.

35 transporting another nucleic acid to which it has been linked. A vector may be characterized by one or a small number of restriction endonuclease sites at which such DNA sequences may be cut in a determinable fashion without the loss of an essential biological function of the vector, and into which a DNA fragment may be spliced in

order to bring about its replication and cloning. A vector may further contain a marker suitable for use in the identification of cells transformed with the vector. Recombinant vectors may be generated to enhance the expression of a gene which has been cloned into it, after transformation into a host. The cloned gene is usually placed under the control of (*i.e.*, operably linked to) certain control sequences or regulatory sequences, which may be either constitutive or inducible.

One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome.

Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". Expression systems for both prokaryotic and eukaryotic cells are described in, for example, chapters 16 and 17 of Sambrook, J. *et al. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.*

In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions. Protocols for producing recombinant retroviruses and for infecting cells *in vitro* or *in vivo* with such viruses can be found in Current Protocols in Molecular Biology, Ausubel, F.M. *et al.* (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines include ψ Crip, ψ Cre, ψ 2 and ψ Am. The genome of adenovirus can be manipulated such that it encodes and expresses a transcriptional regulatory protein but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See for example Berkner *et al.* (1988) *BioTechniques* 6:616; Rosenfeld *et al.* (1991) *Science* 252:431-434; and Rosenfeld *et al.* (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (*e.g.*, Ad2, Ad3, Ad7 *etc.*) are well known to those skilled in the art. Alternatively, an adeno-associated virus vector such as that described in Tratschin *et al.* ((1985) *Mol. Cell. Biol.* 5:3251-3260) can be used.

In general, it may be desirable that an expression vector be capable of replication in the host cell. Heterologous DNA may be integrated into the host genome, and thereafter is replicated as a part of the chromosomal DNA, or it may be DNA which replicates autonomously, as in the case of a plasmid. In the latter case, the vector will
5 include an origin of replication which is functional in the host. In the case of an integrating vector, the vector may include sequences which facilitate integration, *e.g.*, sequences homologous to host sequences, or encoding integrases.

Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are known in the art, and are described in, for example,
10 Powels *et al.* (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985). Mammalian expression vectors may comprise non-transcribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a poly-adenylation
15 site, splice donor and acceptor sites, and transcriptional termination sequences.

The vectors of the subject invention may be transformed into an appropriate cellular host for use in the methods of the invention.

As used interchangeably herein, a "cell" or a "host cell" includes any cultivatable cell that can be modified by the introduction of heterologous DNA. As used herein,
20 "heterologous DNA", a "heterologous gene" or "heterologous polynucleotide sequence" is defined in relation to the cell or organism harboring such a nucleic acid or gene. A heterologous DNA sequence includes a sequence that is not naturally found in the host cell or organism, *e.g.*, a sequence which is native to a cell type or species of organism other than the host cell or organism. Heterologous DNA also includes mutated
25 endogenous genetic sequences, for example, as such sequences are not naturally found in the host cell or organism. Preferably, a host cell is one in which a quorum sensing signal molecule, *e.g.*, an autoinducer molecule, initiates a quorum sensing signaling response which includes the regulation of target quorum sensing controlled genetic sequences. The choice of an appropriate host cell will also be influenced by the choice
30 of detection signal. For example, reporter constructs, as described herein, can provide a selectable or screenable trait upon activation or inhibition of gene transcription in response to a quorum sensing signaling event; in order to achieve optimal selection or screening, the host cell phenotype will be considered.

A host cell of the present invention includes prokaryotic cells and eukaryotic
35 cells. Prokaryotes include gram negative or gram positive organisms, for example, *E. Coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a preferred embodiment, a

host cell of the invention is a mutant strain of *P. aeruginosa* in which *lasI* and *rhlI* are inactivated.

Eukaryotic cells include, but are not limited to, yeast cells, plant cells, fungal cells, insect cells (e.g., baculovirus), mammalian cells, and cells of parasitic organisms, e.g., trypanosomes. Mammalian host cell culture systems include established cell lines such as COS cells, L cells, 3T3 cells, Chinese hamster ovary (CHO) cells, embryonic stem cells, and HeLa cells. Other suitable host cells are known to those skilled in the art.

DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms “transformation” and “transfection” are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

Host cells comprising an isolated nucleic acid molecule of the invention (e.g., a quorum sensing controlled genetic locus operatively linked to a reporter gene) can be used in the methods of the instant invention to identify a modulator of quorum sensing signaling in bacteria.

Exemplification

The invention is further illustrated by the following examples which should not be construed as limiting.

EXAMPLE 1 IDENTIFICATION OF QUORUM SENSING GENES OF *P. AERUGINOSA*

30

Materials and Methods

Bacterial Strains, Plasmids, and Media. The bacterial strains and plasmids used in this example are listed in Table 2.

E. coli and *P. aeruginosa* were routinely grown in Luria-Bertani (LB) broth or LB agar (Sambrook, *et al.* (1989) *Molecular Cloning: a Laboratory Manual*. (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY)), supplemented with antimicrobial agents when necessary. The antimicrobial agents were used at the following concentrations: HgCl₂, 15 µg/ml in agar and 7.5 µg/ml in broth; nalidixic acid 20 µg/ml; carbenicillin, 300 µg/ml; tetracycline, 50 µg/ml for *P. aeruginosa* and 20 µg/ml for *E.*

coli; and gentamicin, 100 µg/ml for *P. aeruginosa* and 15 µg/ml for *E. coli*. Synthetic acyl-HSL concentrations were 2 µM for 3OC₁₂-HSL and 5 µM for C₄-HSL, and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) was used at 50 µg/ml.

5 DNA Manipulations and Plasmid Constructions. DNA treatment with modifying enzymes and restriction endonucleases, ligation of DNA fragments with T4 ligase, and transformation of *E. coli* were performed according to standard methods (Ausubel, F. *et al.* (1997) *Short Protocols in Molecular Biology*. (John Wiley & Sons, Inc., New York, N.Y.)). Plasmid isolation was performed using QIAprep spin miniprep
10 kits (Qiagen Inc.) and DNA fragments were excised and purified from agarose gels using GeneClean spin kits (Bio101 Corp.). DNA was sequenced at the University of Iowa DNA core facility by using standard automated sequencing technology.

To construct pMW10, the pBR322 *tetA(C)* gene-containing *Cla*I-*Not*I DNA fragment in pJPP4 was replaced with a *tetA(B)*-containing *Bst*BI-*Not*I fragment from Tn10. It was necessary to use *tetA(B)* rather than *tetA(C)* to inactivate *lasI* because the *tetA(C)* gene from pBR322 was a hot spot for Tn5::B22 mutagenesis (Berg, D. E. *et al.* (1983) *Genetics* 105, 813-828).

To construct pMW300 a 1.6-kb *Sma*I fragment from pGM Ω 1 that contained the *aacC1* gene (encoding gentamicin acetyltransferase-3-1) was cloned into *Eag*I digested pTL61T, which had been polished with T4 polymerase. The resulting plasmid pTL61T-GM Ω 1 was digested with *Sma*I and *Msc*I to release a 6.5-kb *lacZ-aacC1* fragment. A

TABLE 2. Bacterial strains and plasmids

	Strain or plasmid	Relevant characteristics	Source (reference)
	Strains		
	<i>P. aeruginosa</i> PAO1	Parental strain	(1)
5	<i>P. aeruginosa</i> PDO100	$\Delta rhII::Tn501$ derivative of PAO1, Hg ^r	(2)
	<i>P. aeruginosa</i> PAO-MW1	$\Delta lasI$, $\Delta rhII$ derivative of PDO100, Hg ^r , Tc ^r	This study
	<i>P. aeruginosa</i> PAO-MW10	<i>lasB::lacZ</i> chromosomal insertion in PAO-MW1	This study
	<i>E. coli</i> DH5 α	F ⁻ $\phi 80\Delta lacZ$, $\Delta M15$, $\Delta(lacZYA-argF)U169$, <i>endA1</i> , <i>recA1</i> , <i>hsdR17</i> , <i>deoR</i> , <i>gyrA96</i> , <i>thi-1</i>	(3)
10	<i>E. coli</i> HB101	F ⁻ <i>mcrB</i> , <i>mrr</i> <i>hsdS20</i> , <i>recA13</i> , <i>leuB6</i> , <i>ara-14</i> <i>proA2</i> , <i>lacY1</i> , <i>galK2</i> , <i>xyl-5</i> , <i>mtl-1</i> , <i>rpsL20</i> (Sm ^r), <i>supE44</i>	(3)
	<i>E. coli</i> SY327 λ pir	(λ pir), $\Delta(lac pro)$, <i>argE</i> (Am), <i>rif</i> , <i>nla</i> , <i>recA56</i>	(4)
15	<i>E. coli</i> S17-1	<i>thi</i> , <i>pro</i> , <i>hsdR</i> , <i>recA</i> , <i>RP4-2</i> (Tet::Mu) (Km::Tn7)	(5)
	Plasmids		
	pJPP4	oriR6K, mobRP4, $\Delta lasI$, Tc ^r , Ap ^r	(6)
	pTL61T	<i>lacZ</i> transcriptional fusion vector, Ap ^r	(7)
	pGM Ω 1	Contains <i>aacI</i> flanked by transcriptional and translational stops, Gm ^r	(8)
20	pTL61T-GM Ω 1	pTL61T with <i>aacI</i> gene from pGM Ω 1 upstream of <i>lacZ</i> , Ap ^r , Gm ^r	This study
	pMW100	pJPP4 with 2.7-kb <i>tetA</i> (B) from Tn10 in place of the pBR322 <i>tetA</i> (C), Tc ^r , Ap ^r	This study
25	pRK2013	ori (ColE1), <i>tra</i> ⁺ , (RK2)Km ^r	(9)
	pSUP102	pACYC184 carrying mobRP4, Cm ^r , Tc ^r	(10)
	pSUP102- <i>lasB</i>	pSUP102 carrying <i>lasB</i> on a 3.1-kb <i>P. aeruginosa</i> chromosomal DNA fragment, Cm ^r , Tc ^r	This study
	pMW300	pSUP102- <i>lasB</i> containing <i>lacZ-aacI</i> from pTL61T-GM Ω 1 (<i>lasB-lacZ</i> transcriptional fusion knockout plasmid), Cm ^r , Gm ^r	This study
30	pTn5-B22	pSUP102 with Tn5-B22 (<i>lacZ</i>), Gm ^r	(28)
	Abbreviations for antibiotics are as follows: kanamycin, Km; gentamicin, Gm; ampicillin, Ap; tetracycline, Tc; streptomycin, Sm.		

35

3.1-kb *P. aeruginosa* PAO1 chromosomal DNA fragment containing the *lasB* gene was amplified by PCR using the Expand™ Long Template PCR System (Boehringer Mannheim). This fragment was cloned into *Bam*HI-digested pSUP102. The resulting

plasmid, pSUP102-lasB was digested with *NotI*, polished with T4 polymerase and ligated with the 6.5-kb *lacZ-aacC1* fragment from pTL61T-GM Ω 1 to generate pMW300. The promoterless *lacZ* gene in pMW300 is 549 nucleotides from the start of the *lasB* ORF, it is flanked by 1.5 kb upstream and 1.6 kb downstream *P. aeruginosa* DNA, and it contains the p15A *ori*, which does not support replication in *P. aeruginosa*.

Construction of *P. aeruginosa* Mutants. A *lasI*, *rhlI* mutant strain of *P. aeruginosa* PAO-MW1 was generated by insertional mutagenesis of *lasI* in the *rhlI* deletion mutant, PDO100. For insertional mutagenesis, the *lasI::tetA(B)* plasmid, pMW100 was mobilized from *E. coli* SY327 λ *pir* into PDO100 by triparental mating with the help of *E. coli* HB101 containing pRK2013. Because pMW100 has a λ *pir*-dependent origin of replication, it cannot replicate in *P. aeruginosa*. A tetracycline-resistant, carbenicillin-sensitive exconjugant was selected, which was shown by a Southern blot analysis to contain *lasI::tetA* but not *lasI* or pMW100. To confirm the inactivation of the chromosomal *lasI* in this strain, PAO-MW1, the amount of 3OC₁₂-HSL in the fluid from a stationary phase culture (optical density at 600 nm, 5) was assessed by a standard bioassay (Pearson, J. P. *et al.* (1994) *PNAS*, **91**, 197-201). No detectable 3-OC₁₂-HSL (< 5 nM) was found.

A mutant strain, *P. aeruginosa* PAO-MW10, which contains a *lacZ* reporter in the chromosomal *lasB* gene was constructed by introduction of pMW300 into PAO-MW1 by triparental mating as described above. Exconjugants resistant to gentamicin and sensitive to chloramphenicol were selected as potential recombinants. Southern blotting of chromosomal DNA with *lasB* and *lacZ* probes indicated that the pMW300 *lasB-lacZ* insertion had replaced the wt *lasB* gene.

Southern Blotting. Chromosomal DNA was prepared using the QIAMP tissue kit (Qiagen Inc.). Approximately 2 μ g of chromosomal DNA was digested with restriction endonucleases, separated on a 0.7% agarose gel, and transferred to a nylon membrane according to standard methods (Ausubel, F. *et al.* (1997) *Short Protocols in Molecular Biology*. (John Wiley & Sons, Inc., New York, N.Y.). DNA probes were generated using digoxigenin-11-dUTP by random primed DNA labeling or PCR. The Southern blots were visualized using the Genius™ system as outlined by the manufacturer (Boehringer Mannheim).

Tn5 Mutagenesis. Tn5::B22, which carries a promoterless *lacZ* gene, was used to mutagenize *P. aeruginosa* PAO-MW1 (Simon, R. *et al.* (1989) *Gene* 80, 161-169). Equal volumes of a late logarithmic phase culture of *E. coli* S17-1 carrying pTn5::B22 grown at 30°C with shaking and a late logarithmic phase culture of *P. aeruginosa* PAO-

MW1 grown at 42°C without shaking were mixed. The mixture was centrifuged at 6000 x g for 10 minutes at room temperature, suspended in LB (5% of the original volume), and spread onto LB plates (100 µl per plate). After 16 to 24 hours at 30°C, the cells on each plate were suspended in 500 µl LB and 100 µl volumes were spread onto LB agar plates containing HgCl₂, gentamicin, tetracycline and nalidixic acid. The nalidixic acid prevents growth of *E. coli* but not *P. aeruginosa*. After 48 to 72 hours at 30°C, 20 colonies were selected from each mating and grown on LB selection agar plates containing X-gal. Ten of the 20 were picked for further study. The colonies picked showed a range in the intensity of the blue color on the X-gal plates. In this way, the selection of siblings in a mating were minimized. A Southern blot using a probe to *lacZ* was performed on 20 randomly chosen transconjugants indicated that the Tn5 insertion in each was in a unique location.

The Screen for qsc Fusions. A microtiter dish assay was used to identify mutants showing acyl-HSL-dependent β-galactosidase expression (quorum sensing-controlled or qsc mutants). Each transconjugant was grown in four separate wells containing LB broth without added autoinducer, with added 3OC₁₂-HSL, C₄-HSL, or both 3OC₁₂-HSL and C₄-HSL for 12-16 hours at 37°C. Inocula were 10 µl of an overnight culture and final culture volumes were 70 µl. The β-galactosidase activity of cells in each microtiter dish well was measured in microtiter dishes with a luminescence assay (Tropix) Luminescence was measured with a Lucy I microtiter dish luminometer (Anthos).

Patterns of Acyl-HSL Induction of β-galactosidase Activity in qsc Mutants. The pattern of β-galactosidase expression was examined in response to acyl-HSLs in each of 47 qsc mutants identified in the initial screen. Each mutant was grown in 1 ml of MOPS (50 mM, pH 7.0) buffered LB broth containing one, the other, both, or neither acyl-HSL signal in an 18 mm culture tube at 37°C with shaking. A mid-logarithmic phase culture was used as an inoculum and initial optical densities (ODs) at 600 nm were 0.1. Growth was monitored as OD at 600 nm and β-galactosidase activity was measured in 0.1 ml samples taken at 0, 2, 5, and 9 hours after inoculation.

DNA Sequencing and Sequence Analysis. To identify DNA sequences flanking Tn5::B22 insertions, arbitrary PCR was performed with primers and conditions as described (Caetano-Annoles, G. (1993) *PCR Methods Appl.* 3, 85-92; O'Toole, G. A. et al. (1998) *Mol. Microbiol.* 28, 449-461). Tn5 flanking sequences that could not be identified using arbitrary PCR were cloned. For cloning, 3 µg of chromosomal DNA was digested with *EcoRI* and ligated with *EcoRI*-digested, phosphatase treated pBR322.

E. coli DH5 α was transformed by electroporation with the ligation mixtures and plasmids from gentamicin resistant colonies were used for sequencing Tn5-flanking DNA.

DNA sequences flanking Tn5-B22 insertions were located on the *P. aeruginosa* PAO1 chromosome by searching the chromosomal database at the *P. aeruginosa* Genome Project web site (www.pseudomonas.com). The ORFs containing the insertions are those described at the web site. Functional coupling from the Argonne National Labs (http://wit.mcs.anl.gov/WIT2), sequence analysis, and expression patterns of the qsc mutants were used to identify potential operons (Overbeek, R. *et al.* (1999) *PNAS* 96, 2896-2901).

Results

Identification of *Pseudomonas aeruginosa* qsc Genes. Seven thousand Tn5::B22 mutants of *P. aeruginosa* PAO-MW1 were screened. Tn5::B22 contains a promoterless *lacZ*. *P. aeruginosa* PAO-MW1 is a *lasI*, *rhII* mutant that does not make acyl-HSL signals. Thus, transcription of the Tn5::B22 *lacZ* in a qsc gene was expected to respond to an acyl-HSL signal. The screen involved growth of each mutant in a complex medium in a microtiter dish well with no added acyl-HSL, 3OC₁₂-HSL, C₄-HSL, or both 3OC₁₂-HSL and C₄-HSL. After 12-16 hours, β -galactosidase activity in each culture was measured. Two hundred-seventy mutants showed greater than 2 fold stimulation of β -galactosidase activity in response to either or both acyl-HSL. Of these, 70 showed a greater than 5-fold stimulation of β -galactosidase activity in response to either or both acyl-HSL, and were studied further. Each mutant was grown with shaking in culture tubes and 47 showed a reproducible greater than 5-fold stimulation of β -galactosidase activity in response to either or both of the acyl-HSL signals. These were considered to have Tn5::B22 insertions in qsc genes. It was shown by a Southern blot analysis with a *lacZ* probe that each mutant contained a single Tn5::B22 insertion.

This collection of 47 mutants is not believed to represent the entire set of quorum sensing regulated genes in *P. aeruginosa*. The threshold of greater than 5-fold induction may be too stringent, enough mutants may not have been screened to be assure that insertions in all of the genes in the chromosome have been tested, and there may be conditions other than those which were employed that would have revealed other genes which were not detected in the present screen. Nevertheless, a set of 47 insertions in genes have been identified that show significant activation in response to acyl-HSL (qsc genes).

Responses of qsc Mutants to Acyl-HSL Signals. For cultures of each of the 47 qsc mutants, β -galactosidase activity was measured at different times after addition of

acyl-HSL signals. The basal levels of β -galactosidase varied depending on the mutant. The responses to the acyl-HSL signals could be grouped into 4 general classes based on which of the two signals was required for activation of *lacZ*, and whether the response to the signal(s) occurred immediately or was delayed until stationary phase. A response

5 was considered immediate if there was a 5-fold or greater response within 2 hours of acyl-HSL addition (the optical densities(ODs) of the cultures ranged from 0.5-0.7 at 2 hours). A response was considered delayed or late if there was <5-fold induction at 2 hours but greater than 5-fold induction of β -galactosidase at 5 hours or later (ODs of 2 or greater). In some strains activation of *lacZ* required 3OC₁₂-HSL, others required both

10 3OC₁₂-HSL and C₄-HSL for full activation of *lacZ*. A number of strains responded to either signal alone but showed a much greater response with both 3OC₁₂-HSL and C₄-HSL. None of the mutants responded well to C₄-HSL alone (Table 3). This was expected because expression of RhlR, which is required for a response to C₄-HSL is dependent on 3OC₁₂-HSL (Pesci, E. C. *et al.* (1997) *J. Bacteriol.* 179, 3127-3132).

15 Therefore at least some of the insertions exhibiting a response to both acyl-HSLs may be responding to the *rhl* system, which requires activation by the *las* system.

Class I mutants responded to 3OC₁₂-HSL immediately, Class II responded to 3OC₁₂-HSL late, Class III respond best to both signals early, and Class IV to both signals late. There were 9 Class I, 11 Class II, 18 Class III, and 9 Class IV mutants.

20 Figure 2 shows responses of representative members of each class to acyl-HSLs. Generally, most early genes (Class I and III genes) showed a much greater induction than most late genes (Class II and IV). Many of the Class III mutants showed some response to either signal alone but showed a greater response in the presence of both signals (Table 3 and Figure 2).

25

Identity and Analysis of qsc Genes. The Tn5-B22-marked qsc genes were identified by coupling arbitrary PCR or transposon cloning with DNA sequencing. The sequences were located in the *P. aeruginosa* PAO1 chromosome by searching the *Pseudomonas aeruginosa* Genome Project web site (www.pseudomonas.com). To confirm the

30 locations of the Tn5-B22 insertions in each qsc mutant, a Southern blot analysis was performed with Tn5-B22 as a probe. The sizes of Tn5-B22 restriction fragments were in agreement with those predicted based on the *P. aeruginosa* genomic DNA sequence (data not shown). The 47 qsc mutations mapped in or adjacent to 39 different open reading frames (ORFs). For example Figure 3 depicts the nucleic acid sequence of the

35 quorum sensing controlled locus on the *P. aeruginosa* chromosome mapped in the *P. aeruginosa* mutant strain qsc102.

Table 3. Quorum sensing-controlled genes in *Pseudomonas aeruginosa*

	Classification	Identity ^a	3OC ₁₂ -HSL	Signal response ^b		Genomic Position ^c
				C ₄ -HSL	Both	
5	<u>Class I</u>					
	qsc100	Peptide synthetase	65	3	69	5801998
	qsc101	No match	145	1	184	7730
	qsc102	No match	350	1	400	5547
	qsc103	No match	85	1	95	3961920
10	qsc104	Polyamine binding protein	7	2	8	5402505
	qsc105	FAD-binding protein	40	1	42	5410045
	qsc106A&B	No match	9	1	10	2870317
	qsc107	No match	44	2	50	5799641
15	<u>Class II</u>					
	qsc108	ORF 5	13	1	7	5617382
	qsc109	Bacitracin synthetase 3	13	1	8	5651872
	qsc110A&B	Pyoverdine synthetase D	10	1	7	5661697
	qsc111	Pyoverdine synthetase D	11	1	7	5666282
20	qsc112A&B	Aculeacin A acylase	15	1	12	5701004
	qsc113	Transmembrane protein	5	1	5	3771157
	qsc114 ^c	No match	9	1	7	5209051
	qsc115 ^d	No match	4	1	5	1941557
	qsc116	No match	5	1	5	1138940
25	<u>Class III</u>					
	qsc117 ^d	ACP-like protein	22	22	186	41430
	qsc118	RhlI	38	14	70	4447967
	qsc119	RhlB	9	7	100	4446918
30	qsc120	Chloramphenicol resistance	3	7	24	4592102
	qsc121	3-Oxoacyl ACP synthase	13	27	105	4594988
	qsc122A&B	Cytochrome p450	2	10	90	4595538
	qsc123	9-Cis retinol dehydrogenase	14	28	96	4597340
	qsc124A&B	Pyoverdine synthetase D	35	50	148	4598281
35	qsc125	Zeaxanthin synthesis	20	65	140	4600099
	qsc126	Pristanymycin I synthase 3 & 4	3	5	24	4603518
	qsc127 ^c	No match	5	2	15	4608787
	qsc128	Hydrogen cyanide synthase HcnB	19	12	42	5924799
	qsc129A&B	Cellulose binding protein p40	15	1	100	1141723
40	qsc130	<i>glc</i> operon transcriptional activator	5	1	14	2313744
	qsc131	PhzC	50	168	742	1110
	<u>Class IV</u>					
	qsc132A&B	Unknown (<i>B. pertusis</i>)	1	1	40	3616599
45	qsc133A&B	AcrB	1	1	9	3628342
	qsc134	Saframycin Mx1 synthetase A	6	1	28	3781254
	qsc135	Cytochrome C precursor	3	1	6	4942182
	qsc136 ^c	No match	7	3	45	851491
	qsc137	Asparagine synthetase	1	1	10	2007007
50	qsc138	No match	3	5	32	2459178

^a The bold letters indicate matches were to known *P. aeruginosa* genes.^b The signal response is given as β -galactosidase activity in cells grown in the presence of the indicated signal(s) divided by the β -galactosidase activity of cells grown in the absence of added signals. Maximum responses are indicated.^c The *lacZ* insertions in these strains are in the opposite orientation of the ORFs described in the *P. aeruginosa* Genome Project web site. The insertions are which in locations with no reported identity are been indicated.

^d Insertions do not lie in but are near the putative ORFs indicated. In qsc117 the insertion is 129 bp downstream of the ACP ORF and interrupts a potential rho-independent transcription terminator. The qsc115 insertion is 60 bp upstream of the ORF listed in Materials and Methods.

^e Genomic position as identified using sequence information described in the *P. aeruginosa* Genome Project web site (July 15, 1999 release).

The genomic sequences comprising the ORFs in Table 3 are described in the *Pseudomonas aeruginosa* Genome Sequencing Project web site, as detailed in Table 4.

Only 2 genes were identified that already were known to be controlled by quorum sensing, *rhlI* and *rhlB*. Several other genes potentially involved in processes known to be regulated by quorum sensing were also identified including *phzC* (phenazine synthesis), a putative cyanide synthesis gene (related to the *Pseudomonas fluorescens hcnB*), and ORF 5 (pyoverdine synthesis) (Latifi, A. *et al.* (1995) *Mol. Microbiol.* 17, 333-344; Cunliffe, H. E. *et al.* (1995) *J. Bacteriol.* 177, 2744-2750). Interestingly, *lasB* was not identified by the assay, yet the LasI-LasR quorum sensing system was originally described as regulating *lasB* (Gambello, M. J. *et al.* (1991) *J. Bacteriol.* 173, 3000-3009). A *lasB-lacZ* chromosomal fusion in *P. aeruginosa* PAO-MW1 was constructed, so that regulation of *lasB* by quorum sensing could be compared to the genes identified by the assay. The *lasB-lacZ* fusion only responded slightly to 3OC₁₂-HSL (3-fold stimulation). The full response (12-13-fold over background) required both C₄-HSL and 3OC₁₂-HSL, and the response was late (Figure 2). Thus, *lasB* shows the characteristics of a Class IV gene.

Some of the qsc mutants had obvious phenotypes. Unlike the parent, on LB agar, colonies of the Class II mutants qsc108, 109, 110A and B, and 111 were not fluorescent. Because pyoverdine is a fluorescent pigment, and because the qsc110 and 111 mutations were in genes coding for pyoverdine synthetase-like proteins, it was suspected that these mutations define a region involved in pyoverdine synthesis. The insertion in qsc131 is in *phzC* which is required for pyocyanin synthesis. Although the parent strain produced a blue pigment in LB broth, qsc131 did not. The two qsc132 mutants also did not produce detectable levels of pyocyanin but did produce a water-soluble red pigment.

Functional coupling and sequence analysis were used to identify 7 putative qsc operons, one of which is the previously described *rhlAB* operon (Figure 4). Functional coupling will not organize genes encoding polypeptides without known relatives into operons, and organization of genes in an operon was disallowed in cases where there was greater than 250 bp of intervening sequence between two adjacent ORFs. The

Table 4. ORFs of quorum sensing-controlled genes in *Pseudomonas aeruginosa*

QSC	Insertion July 15, 1999 release	Insertion December 15, 1999 release	Open Reading Frame December 15, 1999 release	Orientation	SEQ ID NO
131	1110	4715256	4714774-4715991	Forward	1
102	5547	2067716	2066736-2068517	Reverse	2
101	7730	2065297	2064803-2065495	Reverse	3
117	41430	2031833	2031245-2031655	Forward	4
136	851491	1221771	1221374-1221961	Reverse	5
116	1138940	934322	934191-935210	Reverse	6
129	1141723	931539	930603-931772	Reverse	7
115	1941557	131753	131583-131792	Reverse	8
137	2007007	66507	66264-68135	Forward	9
130	2313744	6023975	6023787-6024542	Forward	10
138	2459178	5878418	5877776-5878597	Forward	11
106	2870317	5467402	5466520-5467887	Forward	12
132	3616599	4721118	4720249-4721457	Forward	13
133	3628342	4709375	4707483-4710572	Forward	14
113	3771157	4566558	4565369-4567903	Reverse	15
134	3781254	4556461	4555202-4558177	Forward	16
103	3961920	4375793	4375589-4376680	Forward	17
119	4446918	3890793	3890724-3892004	Reverse	18
118	4447967	3889744	3889088-3889738	Reverse	19
120	4592102	3745609	3744850-3746016	Forward	20
121	4594988	3742723	3742643-3743635	Forward	21
122	4595538	3742173	3740961-3742217	Forward	22
123	4597340	3740171	3740054-3740968	Forward	23
124	4598281	3739430	3738724-3740052	Forward	24
125	4600099	3737612	3737561-3738727	Forward	25
126	4603518	3734193	3730455-3737564	Forward	26
127	4608787	3728924		Reverse	
135	4942182	3395532	3395274-3396677	Reverse	27
114	5209051	3128663	3127731-3129116	Forward	28
104	5402505	2935208	2934490-2935593	Forward	29
105	5410045	2927668	2926722-2927972	Reverse	30
108	5617382	2720329	2718890-2720643	Reverse	31
109	5651872	2678258	2671678-2679012	Reverse	32
110	5661697	2676014	2671678-2679012	Reverse	32
111	5666282	2671429	2669119-2671674	Reverse	33
112	5701004	2636707	2636467-2638800	Reverse	34
107	5799641	2538070	2532619-2539008	Reverse	35
100	5801998	2535711	2532619-2539008	Reverse	35
128	5924799	2412909	2412807-2414201	Forward	36

qsc101 and 102 genes are an example of a putative operon that was not identified by
 5 functional coupling (Figure 4). These two ORFs did not show significant similarities
 with other polypeptides. Nevertheless, they are transcribed in the same direction,
 closely juxtaposed, qsc101 and 102 are both Class I genes, and there is a *las* box-like
 element upstream of these ORFs. Expression of the qsc102 insertion is controlled by an

upstream ORF (SEQ ID NO:37) which comprises the sequences between positions 2068711 to 267911 of the *P. aeruginosa* genome (December 15, 1999 release) which in turn is preceded by a *las* box regulatory element (SEQ ID NO:38) which comprises the sequences between positions 2068965 to 2068946 of the *P. aeruginosa* genome (December 15, 1999 release). The *las* box is a palindromic sequence found upstream of and involved in LasR-dependent activation of *lasB* (Rust, L. *et al.*, (1996) *J. Bacteriol.* 178, 1134-1140).

The qsc133A and B insertions are in a putative 3-gene operon with similarity to *acrAB-tolC* from *E. coli* and the *mex-opr* family of efflux pump operons in *P.*

aeruginosa, one of which (*mexAB-oprN*) has been shown to aid 3OC₁₂-HSL efflux (Kohler, T., *et al.* (1997) *Mol. Microbiol.* 23, 345-354; Poole, K., *et al.* (1993) *J. Bacteriol.* 175, 7363-7372; Poole, K., *et al.* (1996) *Mol. Microbiol.* 21, 713-724; Evans, K., *et al.* (1998) *J. Bacteriol.* 180, 5443-5447; Pearson, J. P. *et al.* (1999) *J. Bacteriol.* 181, 1203-1210). The *qsc133* mutations are within a gene encoding a MexF homolog. The *qsc133* mutants show typical Class IV regulation. Expression of *lacZ* is late and dependent on the presence of both acyl-HSL signals (Table 3 and Figure 2). No *las* box-like sequences upstream of this suspected efflux pump operon were identified.

A third possible operon identified by functional coupling is about 8 kb and contains 10 genes. Eight strains with insertions in 6 of the 10 genes were obtained, all of which are Class III mutants (Table 3). A *las* box-like sequence was identified upstream of the first gene of this operon. The function of these 10 genes is unknown but the similarities shown in Table 2 suggest that they may encode functions for synthesis and resistance to an antibiotic-like compound.

The qsc128 mutation is within a gene coding for a polypeptide that shows similarity to the *P. fluorescens hcnB* product and appears to be in a 3-gene operon (Table 3, Figure 4). By analogy to the *P. fluorescens hcn* operon, this operon is likely required for the production of the secondary metabolite, hydrogen cyanide. Previous investigations have shown that hydrogen cyanide production is reduced in *P. aeruginosa rhl* quorum sensing mutants. Consistent with this, qsc128 is a Class III mutant (Table 2). Full induction required both acyl-HSL signals, however, some induction of *lacZ* resulted from the addition of either signal alone (Table 3). A *las* box-like sequence was identified in the region upstream of the translational start codon of the first gene in this operon. This *las*-type box may facilitate an interaction with either LasR or RhlR.

The *phz* operon, required for phenazine biosynthesis, has been described in other pseudomonads and the insertion in strain qsc131 is located in a gene encoding a *phzC* homolog. Analysis of the sequence around this *phzC* homolog revealed an entire phenazine biosynthesis operon, *phzA-G* (Georgakopoulos, D. G. *et al.* (1994) *Appl. Environ. Microbiol.* 60, 2931-2938; Mavrodi, D. V. *et al.* (1998) *J. Bacteriol.* 180,

2541-2548). As discussed above, qsc131 does not produce the blue phenazine pigment pyocyanin. PhzC is part of an operon of several genes including PhzABCDEFG, and transcription of this operon is controlled by the promoter region (SEQ ID NO:39) in front of the first gene in the operon, PhzA. The *phz* operon in *P. aeruginosa* also
 5 contains a *las*-box like sequence upstream of the first gene of the operon. The PhzA promoter region (SEQ ID NO:39) has been cloned into a plasmid, transcriptionally fused to lacZ. The resulting plasmid (pMW303G) was transformed into PAO1 and used as a reporter strain. The resultant bacterial strain generates a quorum sensing signal and responds to it by increased β -galactosidase activity. As shown in Figure 5, this strain
 10 displayed a high level of induction between early and late growth, thus providing a dynamic range for detecting modulation (*e.g.*, inhibition) of quorum sensing signaling. Accordingly this strain may be useful for a single strain assay for identifying for inhibitors of quorum sensing signaling, as described herein.

The final putative operon consists of 2 or 3 genes, qsc109-111, which appear to
 15 be involved in pyoverdine synthesis. These ORFs were not identified in the *P. aeruginosa* genome project web site but were identified and shown to be functionally coupled with the Argonne National Laboratory web site.

For three of the qsc insertions, the *lacZ* gene was in an orientation opposite to the ORF described in the Genome Project web site (qsc114, 127, and 136).
 20

Locations of qsc Genes on the *P. aeruginosa* Chromosome. The qsc genes were mapped to sites on the *P. aeruginosa* chromosome (Figure 6). In addition *lasB*, *lasR* and *lasI*, and *rhlR* were placed on this map. The distribution of currently identified qsc genes is patchy. For example, 16 of the 39 qsc genes representing 3 of the classes
 25 mapped to a 600-kb region of the 6 megabase chromosome. A 140-kb island of 12 Class III genes, 8 transcribed in one direction and 4 transcribed in the other direction (including the *rhl* genes) formed another cluster on the chromosome.

Identification of *las* Box-Like Sequences that Could be Involved in qsc Gene
 30 **Control.** As discussed above, the *las* box is a palindromic sequence found upstream of and involved in LasR-dependent activation of *lasB* (Rust, L. *et al.* (1996) *J. Bacteriol.* 178, 1134-1140). The *las* box shows similarity to the *lux* box, which is the promoter element required for quorum control of the *V. fischeri* luminescence genes (Devine, J. *et al.* (1989) *PNAS* 86, 5688-5692). Elements similar to a *las* box were identified by
 35 searching upstream of qsc ORFs. A search was done for sequences with at least 50% identity to the *las* box found 42 bp upstream of the *lasB* transcriptional start site (Rust, L. *et al.* (1996) *J. Bacteriol.* 178, 1134-1140). *las* box-like sequences were identified which are suspected to be involved in the regulation of 14 of the 39 qsc genes listed in

Table 1 (Figure 7). Because there is little information on the transcription starts of most of the genes identified in the screening assay, some relevant *las* boxes may have been missed and some of the identified sequences may not be in relevant positions.

5 **Discussion**

By screening a library of *lacZ* promoter probes introduced into *P. aeruginosa* PAO1 by transposon mutagenesis, 39 quorum sensing controlled (qsc) genes were identified. Most of these genes were not identified as quorum sensing-controlled previously. Mutations were not found in every gene in putative qsc operons (Figure 4).
 10 Mutants that showed only a small degree of acyl-HSL-dependent *lacZ* induction in the initial screen were not studied. Thus, it is presumed that all of the quorum sensing controlled (qsc) genes have been identified. A conservative estimate is that about 1% of the genes in *P. aeruginosa* are controlled by quorum sensing (39 out of about 5,000-6,000 genes in the *P. aeruginosa* chromosome were confirmed to be qsc without
 15 saturating the mutagenesis). A more liberal estimation of 3-4% can be drawn from the finding of 270 mutants showing at least a 2-fold induction in response to one or both of the acyl-HSL signals in the initial screen of 7,000 mutants.

Several mutants, for example qsc101 and 102 showed an immediate and relatively large response to 3OC₁₂-HSL (Class I mutants, Table 3). The qsc101 and 102
 20 genes code for proteins with no matches in the databases. Several mutants showed a relatively large and immediate response when both signals were supplied in the growth medium. Examples are qsc119 (*rhlB*), 121-125, and 129A and B. The qsc mutant showing the largest response was qsc131. The level of β -galactosidase activity when this mutant was grown in the presence of both signals was greater than 700 times that in the
 25 absence of the signals (Table 3). The qsc131 mutation is in *phzC*, which is a phenazine biosynthesis gene, and the qsc131 mutant did not produce the blue phenazine pigment pyocyanin at detectable levels. Many of the mutants that responded best to both signals early (Class III mutants) showed a small response when exposed to one or the other signal. The reasons for the small response to either signal are unclear at present but the
 30 data suggest that these genes may be subject to signal cross talk, or they may show a response to either LasR or RhlR. One reason they may respond to both signals better than they respond to C₄-HSL alone is that 3OC₁₂-HSL and LasR are required to activate RhlR, the transcription factor required for a response to C₄-HSL (Latifi, A. *et al.* (1996) *Mol. Microbiol.* 21, 1137-1146; Pesci, E. C. *et al.* (1997) *J. Bacteriol.* 179, 3127-3132).
 35 There were two mutant classes that showed a delayed response to the signals; Class II mutants which required only 3OC₁₂-HSL, and Class IV mutants, which required both signals for full induction. These mutants showed between 5 and 45-fold activation of gene expression (Table 3). There are a number of possible explanations for a delayed

response to signal addition. It is possible that some of these genes are stationary phase genes. It is also possible that some are iron repressed. For example, it is known that the synthesis of pyoverdine is regulated by iron and the Class II, delayed response, qsc108-111 mutations are in genes involved in pyoverdine synthesis (Cunliffe, H. E. *et al.*

5 (1995) *J. Bacteriol.* 177, 2744-2750; Rombel, I. *et al.* (1995) *Mol. Gen. Genet.* 246, 519-528). It is also possible that some of these genes are not regulated by quorum sensing, directly. The acyl-HSL signals might control other factors that influence expression of any of the genes that have been identified and this possibility seems most likely with the late genes in Classes II and IV. Indirect regulation may not be common
10 for late genes. This is known because the *lasB-lacZ* chromosomal insertion which was generated by site-specific mutation was in Class IV, and it is known from other investigations that *lasB* responds to LasR and 3OC₁₂-HSL, directly (Passador, L. *et al.* (1993) *Science* 260, 1127-1130; Rust, L. *et al.* (1996) *J. Bacteriol.* 178, 1134-1140). The two classes of late qsc genes may be comprised of several subclasses.

15 *Las* boxes are genetic elements which may be involved in the regulation of qsc genes. Although sequences with characteristics similar to *las* boxes were identified, (Figure 7), the locations of these sequences have not provided insights about the differences in the patterns of gene expression among the four classes of genes. It is possible that when the promoter regions of the qsc genes are studied that particular
20 motifs in the regulatory DNA of different classes of genes will be revealed.

Many of the qsc genes appear to be organized in two patches or islands on the *P. aeruginosa* chromosome (Figure 7). Because LasR mutants are defective in virulence it is tempting to speculate that these gene clusters may represent pathogenicity islands. The *rhII-rhIR* quorum sensing modulation occurs on one of the qsc islands, but none of
25 the qsc genes are tightly linked to the *lasR-lasI* modulon. Genes representing each of the 4 classes occur over the length of the chromosome and on both DNA strands. This is consistent with the view that quorum sensing is a global regulatory system in *P. aeruginosa*. Of interest there is a third LuxR family member in *P. aeruginosa*. This gene is adjacent to and divergently oriented from qsc103.

30 Quorum sensing is critical for virulence of *P. aeruginosa* and for the development of mature biofilms. The methodology disclosed herein for identification of qsc genes provides a manageable group of genes to test for function in virulence and biofilm development. Furthermore, the availability of the *P. aeruginosa* genome sequence will very likely lead to the development of a gene expression microarray for
35 this organism. The methods described herein provide a set of 39 genes that respond to specific treatments in a predictable fashion (Table 3).

EXAMPLE 2 SCREENING ASSAY FOR QUORUM SENSING INHIBITING COMPOUNDS


In this example, the screening assay used two strains of *P. aeruginosa*: a wild type *P. aeruginosa* (PAO1) and QSC102, from Example 1 (see Figure 8). This assay will detect inhibition of all aspects of quorum sensing signaling, *e.g.*, signal generation and signal reception.

Procedural Overview


Microtiter plates are prepared by adding 200 μ L Luria Broth ("LB") agar, containing 0.008 % 5-bromo-4-chloro-3-indolyl- β -D-galactose (X-gal) to each well. Overnight cultures of PAO1 and QSC102 are subcultured in LB to a starting absorbance at 600 nm ("A600") of 0.05 and grown at 37 $^{\circ}$ C to an A600 of 1.0. PAO1 is diluted 2.5 x 10⁵-fold in LB and 5 μ L of this is applied to the surface of the LB agar in each well. Plates are then dried in a laminar flow hood for 60 minutes. A tenfold dilution of QSC102 in LB is used to inoculate each well using a replicator. Plates are then sealed and incubated at 37 $^{\circ}$ C for 40 hours. Growth and color development are evaluated visually and the data is recorded with a camera.

The test compound was present in a microtiter well and overlaid with LB agar and 5-bromo-4-chloro-3-indolyl- β -D-galactose (X-gal). Both strains were spotted on the agar in each well. PAO1 emitted the acyl-HSL signal (3-oxo-C12-HSL), to which QSC102 responded by turning blue. QSC102 expressed β -galactosidase only in response to the LasI signal (3-oxo-C12-HSL); the *lacZ* fusion in QSC102 did not respond to the RhlI signal (C4-HSL). Hence, the assay was selective for inhibitors of the Las system. Inhibition of signaling was evaluated qualitatively by the absence or weakening of the blue color development.

The assay was used to test 6 product analogs, two of which showed an inhibitory effect: butyrolactone and acetyl-butyrolactone. Although bacterial growth was not inhibited, the color development was reduced. Color reduction correlated directly with test compound concentration, although relatively high concentrations (~20 mM) were required to suppress color development completely (Figure 9).



butyrolactone



acetyl-butyrolactone

EXAMPLE 3 DEVELOPMENT OF A *P. AERUGINOSA* STRAIN FOR A HIGH THROUGHPUT SCREENING ASSAY

A. Construction of Reporter Strain-Chromosomal Insertion of Reporter

A strain for use in high-throughput screening was constructed by inserting the lacZ transcriptional fusion, linked gentamicin resistance marker, and about 2 kb of flanking DNA from strain QSC102 into a mobilizable plasmid (such as pSUP102) as depicted in Figure 10A. Plasmid pSUP102 confers tetracycline resistance and does not replicate in *P. aeruginosa* (Simon, R. *et al.* (1986) *Meth. Enzym.* 118:640-659). The pSUP102-derivative was then transferred into PAO1 by bi- or triparental mating, selecting for gentamicin resistance (Suh, S. J. *et al.* (1999) *J Bacteriol.* 181(13):3890-7). Gentamicin resistant isolates were screened for tetracycline sensitivity (*i.e.*, a double cross-over event has resulted in a chromosomal insertion). Southern blotting was used to confirm the nature of the recombination event and to rule out candidates with more than one insertion. The resultant bacterial strain generates the signal (3-oxo-C12-HSL) and responds to it by increased β -galactosidase activity. A similar strategy is used to create a reporter strain that expresses *gfp* instead of *lacZ*. The initial GFP variant is the stable and bright variant GFPmut2 (Cormack, B. P. *et al.* (1996) *Gene.* 173(1):33-38).

Procedural Overview of Assay

A culture of PAQ1 reporter strain (carrying the reporter gene lacZ transcriptionally fused to the regulatory sequence of qsc102 in the wildtype background, PAO1) was grown in LB, 100 μ g/ml gentamicin overnight, such that the A600 was around 0.1. The culture was washed in LB twice and used to subculture at a 1:1000 dilution in LB. The subculture was grown in the presence or absence of test compound. Growth was monitored at A600 and expression of β -galactosidase activity is measured according to the Miller assay (Miller, J. A. (1976) in *Experiments in Molecular Genetics* pp 352-355, Cold Spring Harbor Lab. Press, Plainview, NY).

The reporter strain was tested by growing it in microtiter plates in the presence and absence of known inhibitors of bacterial signaling. Examples of known inhibitors are: acetyl-butyrolactone, butyrolactone, and methylthioadenosine, a product of the synthase reaction that was shown to be inhibitory to the RhlI synthase (Parsek, M. R. *et al.* (1999) *Proc. Natl. Acad. Sci. USA.* 96:4360-4365). Initial characterization of the assay entailed following the optical density (cell growth) in individual sample wells and measuring induction levels at different time points. Figure 10B shows the induction of β -galactosidase as PAQ1 reaches high density, wherein cell growth is measured at 600

nm (closed circles) and expression of β -galactosidase is measured in Miller units (open circles). For GFP fusions, the fluorescence of the culture is determined after excitation at 488 nm.

5 **B. Construction of Reporter Strain-Reporter on a Plasmid**

The PAO1/pMW303G strain is constructed as described in Example 1 above.

Procedural Overview of the Assay

10 An overnight culture of PAO1/pMW303G was diluted to an A600 of 0.1 in LB, 300 μ g/ml carbenicillin. Of this, 50 μ L were added to microtiter plate wells and grown at 37 °C, shaking at 250 rpm, in the presence or absence of test compounds. Culture growth was monitored directly in the microtiter plate at 620 nm. Expression of the reporter gene, β -galactosidase was measured with the Galacton substrate by Tropix as follows. 12A 20 μ L aliquot of the culture was added to 70 μ L of 1:100 diluted Galacton
15 substrate (Tropix, PE Biosystems, Bedford, MA) and incubated in the dark at room temperature for 60 minutes. The reaction was stopped and light emission was triggered by the addition of 100 μ L Accelerator II (Tropix, PE Biosystems, Bedford, MA), and luminescence was read with plate reader (SpectrofluorPlus, Tecan). Timepoints were taken at 5, 8 and 12 minutes.

20 In either embodiment of the assay (chromosomal insertion of reporter, or reporter on a plasmid), a satisfactory assay shows normal cell growth but reduced β -galactosidase activity or *gfp* expression in the presence of a known signaling inhibitor. Possible problems associated with the use of fluorescence in whole-cell systems are interference by turbidity as cell density increases and the production of pyocyanin and
25 pyoverdine, fluorescent molecules that are excreted by wild type *P. aeruginosa*. However, interference due to endogenous fluorescent pigments may be reduced by using mutants that lack these pigments (Byng, G. S. *et al.* (1979) *J Bacteriol.* 138(3):846-52).

30 **EXAMPLE 4 SCREENING ASSAY TO DETERMINE INHIBITION OF THE SIGNAL SYNTHASE**

 An assay was developed to measure inhibition of RhII activity, based on a previously published enzyme assay for RhII (Parsek, M. R. *et al.* (1999) *Proc. Natl.*
35 *Acad. Sci. USA.* 96:4360-4365). It was shown that the substrates for RhII are S-adenosylmethionine (SAM) and butanoyl-acyl carrier protein (C4-ACP). It is proposed that RhII can be used as a model enzyme to study inhibition of acyl-HSL synthases. This is based on the observation that TraI from *Agrobacterium tumefaciens* (Moré, M. I. *et al.* (1996) *Science.* 272(5268):1655-8) and LuxI from *Vibrio fischeri* (Schaefer, A. L.

et al. (1996) *Proc Natl Acad Sci U S A.* 93(18):9505-9), two homologs of RhlI and LasI, that also utilize SAM and the respective acylated-acyl carrier protein as their substrates.

RhlI activity assay. Studies of autoinducer synthases have been hampered by the low solubility of the enzyme. It is only in the past year that the first rigorous characterization of an autoinducer synthase was published (Parsek, M. R. *et al.* (1999) *Proc. Natl. Acad. Sci. USA.* 96:4360-4365). This study was performed on RhlI, which had been slightly overproduced in a LasI minus strain of *P. aeruginosa*, thereby avoiding previously encountered problems of solubility. The reaction mechanism deduced for RhlI is summarized in Figure 11. The substrates for the synthase are butanoyl-acyl carrier protein (C4-ACP) and S-adenosylmethionine (SAM). The amino-group of SAM attacks the thioester of C4-ACP to form a peptide bond between butanoic acid and SAM. The first product, acyl carrier protein (ACP) is released. Next, the SAM-moiety undergoes internal ring closure to form a homoserine lactone (HSL). Methylthioadenosine (MTA) and butanoyl-HSL (C4-HSL) are released.

The enzyme assay reaction mixture contains 60 μM ^{14}C -labeled SAM and 40 μM C4-ACP in a final volume of 100 μL (buffer: 2 mM dithiothreitol, 200 mM NaCl, 20 mM Tris-HCL, pH 7.8). The reaction is started with the addition of 70 ng RhlI, incubated at 37 °C and quenched after 10 min by addition of 4 μL of 1 M HCl. Product formation is quantitated by extracting the reaction mixtures with 100 μL ethyl acetate and scintillation counting the radiolabeled C4-HSL, which partitions into the organic phase. (SAM remains in the aqueous phase.)


Other variations on the assay include detection of the non-acylated ACP (i.e., ACP with a free thiol group). Non-acylated ACP can be detected through the use of a thiol reagent such as dithionitrobenzoic acid (DTNB), which releases a highly colored thiolate ($\epsilon_{412} = 13\,600\text{ cm}^{-1}\text{ M}^{-1}$) upon reaction with thiol groups (Ellman, G. L. (1959) *Arch. Biochem. Biophys.* 82:70-77). Another variation of this assay uses an even more sensitive reagent, 4,4'-dithiobipyridyl which has a $\epsilon_{324} = 20\,000\text{ cm}^{-1}\text{ M}^{-1}$ (Jamin, M. *et al.* (1991) *Biochem J.* 280(Pt 2):499-506). Use of DTNB eliminates the need for radioactivity and allows for a continuous assay.

Another variation on the assay includes using a substitute for the substrate C4-ACP. It has already been found that RhlI turns over butanoyl-CoA in lieu of C4-ACP (Parsek, M. R. *et al.* (1999) *Proc. Natl. Acad. Sci. USA.* 96:4360-4365). The K_M for the CoA substrate is 230 μM , compared to 6 μM for C4-ACP, but v_{max} is only one order of magnitude slower. N-Acetylcysteamine represents a truncated moiety of CoA and acylated N-acetylcysteamines often function as substrate analogs for CoA-dependent enzymes (Bayer *et al.* (1995) *Arch Microbiol.* 163(4):310-2; Singh, N. *et al.* (1985) *Biochem Biophys Res Commun.* 131(2):786-92; Whitty, A. (1995) *Biochemistry*.

004050 "0645360

34(37):11678-89). It will be determined whether butanoyl-N-acetylcysteamine is turned over by RhII. If so, an assay will be developed for the release of free thiol groups with a thiol reagent such as DTNB. Butanoyl-N-acetylcysteamine is readily synthesized from the commercially available precursors butyrylchloride and N-acetylcysteamine.

5



butanoyl-N-acetylcysteamine

LasI activity assay. In analogy with RhII, TraI, and LuxI, proposed substrates for LasI are SAM and 3-oxo-C12-ACP. In this assay, compounds are tested for inhibiting the activity of LasI. This assay is based on observations that bacterial strains incubated with ^{14}C -labeled methionine produce radiolabeled acylated-HSLs, which can be isolated from the culture supernatant and identified by their retention times (in comparison to known standards) when eluted over a high pressure liquid chromatography (HPLC) reversed phase column. A synthase-inhibitor assay has been set up using this methodology.

A *Pseudomonas* strain that expresses *lasI* but not *rhII*, such as PDO100, is grown in the presence and absence of the test compound (Brint, J. M. *et al.* (1995) *J Bacteriol.* 177(24):7155-63). Cells are pulsed for 10-30 minutes with ¹⁴C-labeled methionine (available from American Radiochemicals) and pelleted by centrifugation. The supernatant liquid is extracted with ethyl acetate and the products separated by HPLC. If the test compound inhibits LasI synthase, the amount of 3-oxo-C12-HSL produced will be significantly reduced when compared to the control.

An *in vitro* assay for LasI activity similar to the radiometric assay used to study RhlI will be developed. The substrates for this assay are ^{14}C -labeled SAM (available Amersham Pharmacia) and 3-oxo-C12-ACP (similar methodology in Moré, M. I. *et al.* (1996) *Science*. 272(5268):1655-8). LasI activity is monitored by the appearance of radiolabeled 3-oxo-C12-HSL, after extraction into ethyl acetate and scintillation counting. Initially, crude extracts of LasI overexpressed in *E. coli* serve as the source of enzyme. Once a satisfactory assay is in place, a purification protocol will be developed to obtain LasI in a soluble and active form. The purification may involve expression at low levels (low plasmid copy number, weak promoter, low growth temperature) in a *P. aeruginosa rhlI* mutant. Purification will follow standard techniques such as ammonium sulfate precipitation, anion exchange chromatography, cation exchange chromatography and size-exclusion chromatography.

**EXAMPLE 5 IN VIVO ASSAYS TO DETERMINE INHIBITION OF
SIGNAL BINDING**

5

In vivo assays were also used to determine whether a test compound inhibits signal reception by LasR.

One assay used the *P. aeruginosa* strain QSC102 (Table 3), which responds to the presence of exogenous 3-oxo-C12-HSL by inducing β -galactosidase activity up to
10 400-fold (Example 1). Cells were grown in the presence of a minimal concentration of 3-oxo-C12-HSL and in the presence and absence of the test compound. If the test compound interferes with signal reception, β -galactosidase activity is reduced. Interference can be a result of any of several mechanisms. The simplest is, if the test compound prevents the 3-oxo-C12-HSL from binding to LasR. Alternatively, the test
15 compound may prevent LasR from binding to DNA or interacting productively with RNA polymerase.

A further *in vivo* assay is used to determine whether a test compound inhibits binding of 3-oxo-C12-HSL to LasR. This assay is based on an observation originally made with LuxR of *Vibrio fischeri*. Namely, the autoinducer binds to *Escherichia coli*
20 cells in which LuxR is produced, provided that LuxR is co-expressed with Hsp60 (Adar *et al.* (1993) *J Biolumin Chemilumin.* 8(5):261-6). This finding was used to develop a competition-assay for binding of inhibitors to LuxR (Schaefer, A. L. *et al.* (1996) *J Bacteriol.* 178(10):2897-901) and LasR (Passador, L. *et al.* (1996) *J Bacteriol.* 178(20):5995-6000). Briefly, cultures of *E. coli* harboring expression plasmids for
25 Hsp60 and LasR (or LuxR) are induced for several hours, at which time an aliquot of cells is added to tritiated signal molecule, alone or in combination with a potential inhibitor. After 10-15 minutes, cells are pelleted by centrifugation, washed, and the amount of radioactivity bound to the cells is determined by scintillation counting.

Plasmids for expression of LasR (pKDT37) (Passador, L. *et al.* (1996) *J*
30 *Bacteriol.* 178(20):5995-6000) and Hsp60 (pGroESL) have been made. A simple method for preparing ^{14}C -labeled 3-oxo-C12-HSL has been developed. *E. coli* cells expressing *lasI* excrete ^{14}C -labeled 3-oxo-C12-HSL into the medium when incubated in the presence of ^{14}C -labeled methionine. The ^{14}C -labeled 3-oxo-C12-HSL can be recovered by extraction into ethyl acetate and purified by HPLC. The correct product is
35 identified by its radioactivity and by the correct HPLC retention time compared to an unlabeled standard.

EXAMPLE 6 ASSAY FOR INHIBITION OF BIOFILMS

This assay tests whether compounds useful for inhibiting quorum sensing also inhibit or modulate the formation or growth of biofilms. The LasI/LasR signaling system was found to regulate not only the expression of virulence factors, but also the development of mature biofilms (Davies, D. G. *et al.* (1998) *Science*. 280(5361):295-8). This was demonstrated by using a simple flow-through system, as shown in Figure 12, that allows fresh medium to be pumped through a small chamber in a Plexiglas body.

Cultures of *P. aeruginosa* expressing green fluorescent protein (GFP) were grown in a chamber that was sealed with a coverslip and flushed with fresh medium. Surface attachment and biofilm maturation were determined by examining the coverslip by epifluorescence and confocal microscopy. Both wild type PAO1 and a *rhlI* mutant strain were able to attach to the surface and form the mushroom-shaped structure characteristic of a biofilm. However, a *lasI* mutant that cannot synthesize the signal molecule 3-oxo-C12-HSL was only able to attach to the surface. It did not encase itself in an extracellular matrix or form any kind of three-dimensional structure. It also remained susceptible to 0.2 % sodium dodecyl sulfate, which was used to mimic the susceptibility to a biocide. When the 3-oxo-C12-HSL signal was added back to the *lasI* mutant cells, the wild type phenotype was restored. The cells formed biofilms and remained resistant to sodium dodecyl sulfate.

Accordingly, the bioreactor depicted in Figure 12 is inoculated with wild type *P. aeruginosa* PAO1 that expresses GFP. Test compounds (signaling inhibitors) are added to the flow-through medium to determine whether they prevent formation of the three-dimensional structures typical of a bacterial biofilm. Biofilm formation is monitored using a confocal microscope.

References

1. Holloway, B. W., Krishnapillai, V. & Morgan, A. F. (1979) *Microbiol. Rev.* **43**, 73-102.
2. Brint, J. M. & Ohman, D. E. (1995) *J. Bacteriol.* **177**, 7155-7163.
3. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular cloning: a laboratory manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).
4. Miller, V. L. & Mekalonos, J. J. (1988) *J. Bacteriol.* **170**, 2575-2583.

5. Simon, R., Prierer, U. & Puhler, A. (1983) *Bio-Technology* **1**, 37-45.
6. Pearson, J. P., Pesci, E. C. & Iglewski, B. H. (1997) *J. Bacteriol.* **179**, 5756-5767.
7. Linn, T. & St Pierre, R. (1990) *J. Bacteriol.* **172**, 1077-1084.
- 5 8. Schweizer, H. P. (1993) *Biotechniques* **15**, 831-833.
9. Figurski, D. H. & Helinski, D. R. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 1648-1652.
10. Simon, R., O'Connell, M., Labes, M. & Puhler, A. (1986) in *Methods in Enzymology*, Vol. 118, pp. 640-659.

Incorporation by Reference

The contents of all references, patents and published patent applications cited throughout this application, as well as the figures and the sequence listing, are incorporated herein by reference.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments and methods described herein. Such equivalents are intended to be encompassed by the scope of the following claims.

What is claimed is:

✓ 1. A method for identifying a modulator of quorum sensing signaling in bacteria, said method comprising:

5

providing a cell which comprises a quorum sensing controlled gene, wherein said cell is responsive to a quorum sensing signal molecule such that a detectable signal is generated;

10 contacting said cell with a quorum sensing signal molecule in the presence and absence of a test compound;

and detecting a change in the detectable signal to thereby identify said test compound as a modulator of quorum sensing signaling in bacteria.

15

2. The method of claim 1, wherein said cell further comprises means for generating said detectable signal.

20 3. The method of claim 2, wherein said signal generation means comprises a reporter gene, and wherein said quorum sensing signal molecule causes transcription of said reporter gene, said transcription providing said detectable signal.

25 4. The method of claim 3, wherein said reporter gene is operatively linked to a regulatory sequence of said quorum sensing controlled gene.

5. The method of claim 4, wherein said reporter gene is selected from the group consisting of *ADE1*, *ADE2*, *ADE3*, *ADE4*, *ADE5*, *ADE7*, *ADE8*, *ASP3*, *ARG1*, *ARG3*, *ARG4*, *ARG5*, *ARG6*, *ARG8*, *ARO2*, *ARO7*, *BAR1*, *CAT*, *CHO1*, *CYS3*, *GAL1*, *GAL7*, *GAL10*, *GFP*, *HIS1*, *HIS3*, *HIS4*, *HIS5*, *HOM3*, *HOM6*, *ILV1*, *ILV2*, *ILV5*, *INO1*, *INO2*, *INO4*, *lacZ*, *LEU1*, *LEU2*, *LEU4*, *luciferase*, *LYS2*, *MAL*, *MEL*, *MET2*, *MET3*, *MET4*, *MET8*, *MET9*, *MET14*, *MET16*, *MET19*, *OLE1*, *PHO5*, *PRO1*, *PRO3*, *THR1*, *THR4*, *TRP1*, *TRP2*, *TRP3*, *TRP4*, *TRP5*, *URA1*, *URA2*, *URA3*, *URA4*, *URA5* and *URA10*.

35 6. The method of claim 5, wherein said reporter gene is *lacZ* or *GFP*.

7. The method of claim 1, wherein said cell does not express said quorum sensing signal molecule.

8. The method of claim 7, wherein said quorum sensing signal molecule is produced by a second cell.

5 9. The method of claim 1, wherein said cell is a prokaryote or eukaryote.

10. The method of claim 9, wherein said cell is a bacterium.

11. The method of claim 8, wherein said second cell is a prokaryote or
10 eukaryote.

12. The method of claim 11, wherein said second cell is a bacterium.

13. The method of claims 10 or 12, wherein said bacterium is a gram
negative bacterium.

15 14. The method of claim 13, wherein said gram negative bacterium is
Pseudomonas aeruginosa.

20 15. The method of claim 10, wherein said bacterium is a mutant strain of
Pseudomonas aeruginosa which comprises a regulatory sequence of a quorum sensing
controlled gene operatively linked to a reporter gene, wherein in said mutant strain, *lasI*
and *rhlI* are inactivated.

25 16. The method of claim 12, wherein said second cell is wild type
Pseudomonas aeruginosa.

17. The method of claim 1, wherein said quorum sensing controlled gene is
endogenous to said cell.

30 18. The method of claim 10, wherein said quorum sensing controlled gene
encodes a virulence factor.

19. The method of claim 10, wherein said quorum sensing controlled gene
encodes a polypeptide which inhibits a bacterial host defense mechanism.

35 20. The method of claim 10, wherein said quorum sensing controlled gene
encodes a polypeptide which regulates biofilm formation.

21. The method of claim 1, wherein said quorum sensing signal molecule is an autoinducer of said quorum sensing controlled gene.

22. The method of claim 21, wherein said autoinducer is a homoserine
5 lactone.

23. The method of claim 22, wherein said test compound is a homoserine lactone analog.

10 24. The method of claim 1, wherein said modulator inhibits an enzyme involved in the synthesis by said bacterium of said quorum sensing signal molecule.

25. The method of claim 1, wherein said modulator inhibits reception of said quorum sensing signal molecule by said bacterium.

15 26. The method of claim 1, wherein said modulator scavenges said quorum sensing signal molecule.

27. A method for identifying a modulator of quorum sensing signaling in
20 *Pseudomonas aeruginosa*, said method comprising:

providing a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule;

25 providing a mutant strain of *Pseudomonas aeruginosa* which comprises a reporter gene operatively linked to a regulatory sequence of a quorum sensing controlled gene, wherein said mutant strain is responsive to said quorum sensing signal molecule produced by said wild type strain, such that a detectable signal is generated;

30 contacting said mutant strain with said quorum sensing signal molecule and a test compound; and

detecting a change in the detectable signal to thereby identify said test compound as a modulator of quorum sensing signaling in *Pseudomonas aeruginosa*.

35

28. The method of claim 27, wherein in said mutant strain, *lasI* and *rhII* are inactivated.

29. The method of claim 27, wherein said reporter gene is *lacZ* or *GFP*.

30. The method of claim 29, wherein said reporter gene is *lacZ*.

5 31. The method of claim 29, wherein said reporter gene is *GFP*.

32. The method of claim 31, wherein said reporter gene is a variant of *GFP*.

33. The method of claim 32, wherein said variant is GFPmut2.

10

34. The method of claim 27, wherein said mutant strain of *Pseudomonas aeruginosa* comprises a promoterless reporter gene inserted at a genetic locus in the chromosome of said *Pseudomonas aeruginosa*, wherein said locus comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36.

35. The method of claim 34, wherein said promoterless reporter gene is inserted in said chromosome at a locus comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:28 and SEQ ID NO:35.

36. The method of claim 34, wherein said reporter gene is contained in a transposable element.

30

37. A mutant strain of *Pseudomonas aeruginosa* comprising a promoterless reporter gene inserted at a genetic locus in the chromosome of said *Pseudomonas aeruginosa*, wherein said locus comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID

35

NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36.

5 38. The mutant strain of claim 37, wherein said reporter gene is contained in a transposable element.

 39. The mutant strain of claim 37, wherein *lasI* and *rhlI* are inactivated.

10 40. The mutant strain of claim 37, wherein said strain is responsive to a quorum sensing signal molecule such that a detectable signal is generated by said reporter gene.

 41. The mutant strain of claim 37, wherein said reporter gene is *lacZ* or *GFP*.

15 42. The method of claim 41, wherein said reporter gene is a variant of *GFP*.

 43. The method of claim 42, wherein said variant is GFPmut2.

20 44. A method for identifying a modulator of a quorum sensing signaling in *Pseudomonas aeruginosa*, said method comprising:

 providing a wild type strain of *Pseudomonas aeruginosa* which produces a quorum sensing signal molecule;

25 providing a mutant strain of *Pseudomonas aeruginosa* which comprises a promoterless reporter gene inserted at a genetic locus in the chromosome of said *Pseudomonas aeruginosa*, wherein said locus comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID
30 NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID
35 NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36; and wherein said mutant strain is responsive to said quorum sensing signal molecule produced by said wild type strain, such that a detectable signal is generated by said reporter gene;

contacting said mutant strain with said quorum sensing signal molecule and a test compound; and

- 5 detecting a change in the detectable signal to thereby identify said test compound as a modulator of quorum sensing signaling in *Pseudomonas aeruginosa*.

45. The method of claim 44, wherein said reporter gene is contained in a transposable element.

10

46. An isolated nucleic acid molecule comprising a nucleotide sequence, said nucleotide sequence comprising:

15 a regulatory sequence derived from the genome of *Pseudomonas aeruginosa*, wherein said regulatory sequence regulates a quorum sensing controlled genetic locus of the *Pseudomonas aeruginosa* chromosome, and wherein said locus comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36; and

25

a reporter gene operatively linked to said regulatory sequence.

47. An isolated nucleic acid molecule comprising a quorum sensing controlled genetic locus derived from the genome of *Pseudomonas aeruginosa*, wherein
30 said locus comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID

35

NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36,
operatively linked to a reporter gene.

48. An isolated nucleic acid molecule comprising a polynucleotide having at
least 80% identity to a quorum sensing controlled genetic locus derived from the
genome of *Pseudomonas aeruginosa*, wherein said locus comprises a nucleotide
sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID
NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ
ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID
NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID
NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID
NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID
NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID
NO:34, SEQ ID NO:35 and SEQ ID NO:36, operatively linked to a reporter gene.

49. An isolated nucleic acid molecule comprising a polynucleotide that
hybridizes under stringent conditions to the complement of a nucleotide sequence
comprising a quorum sensing controlled genetic locus derived from the genome of
Pseudomonas aeruginosa, wherein said locus comprises a nucleotide sequence selected
from the group consisting of: SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID
NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ
ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID
NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID
NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID
NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID
NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID
NO:35 and SEQ ID NO:36, operatively linked to a reporter gene.

50. The nucleic acid molecule of any one of claims 46, 47, 48 and 49,
wherein said reporter gene is contained in a transposable element.

51. A vector comprising the isolated nucleic acid molecule of any one of
claims 46, 47, 48 and 49.

52. A cell containing an isolated nucleic acid molecule of any one of claims 46, 47, 48 and 49.

53. A method for identifying a modulator of quorum sensing signaling in
5 bacteria, said method comprising:

providing the cell of claim 52, wherein said cell is responsive to a quorum sensing signal molecule such that a detectable signal is generated;

10 contacting said cell with a quorum sensing signal molecule in the presence and
absence of a test compound;

and detecting a change in the detectable signal to thereby identify said test compound as a modulator of quorum sensing signaling in bacteria.

54. A compound which inhibits quorum sensing signaling in *Pseudomonas aeruginosa*, said compound having been identified by the method of claim 28.

55. The compound of claim 54, which inhibits quorum sensing signaling in
20 *Pseudomonas aeruginosa* by inhibiting an enzyme involved in the synthesis of a quorum
sensing signal molecule, by interfering with quorum sensing signal reception , or by
scavenging the quorum sensing signal molecule.

56. A method for identifying a quorum sensing controlled gene in bacteria,
25 said method comprising:

providing a cell which is responsive to a quorum sensing signal molecule such that expression of a quorum sensing controlled gene is modulated, and wherein modulation of the expression of said quorum sensing controlled gene generates a detectable signal;

contacting said cell with a quorum sensing signal molecule;

and detecting a change in the detectable signal to thereby identify a quorum
35 sensing signaling controlled gene in bacteria.

57. The method of claim 56, wherein said cell further comprises means for generating said detectable signal.

58. The method of claim 57, wherein said signal generation means comprises a reporter gene, and wherein modulation of the expression of said quorum sensing controlled gene modulates transcription of said reporter gene, said transcription
5 providing said detectable signal.

59. The method of claim 58, wherein said reporter gene is operatively linked to a regulatory sequence of said quorum sensing controlled gene.

10 60. The method of claim 58, wherein said reporter gene is opertively linked to said quorum sensing controlled gene.

61. The method of either of claims 59 and 60, wherein said reporter gene is contained in a transposable element.
15

62. The method of claim 58, wherein said reporter gene is selected from the group consisting of *ADE1*, *ADE2*, *ADE3*, *ADE4*, *ADE5*, *ADE7*, *ADE8*, *ASP3*, *ARG1*, *ARG3*, *ARG4*, *ARG5*, *ARG6*, *ARG8*, *ARO2*, *ARO7*, *BARI*, *CAT*, *CHO1*, *CYS3*, *GAL1*, *GAL7*, *GAL10*, *GFP*, *HIS1*, *HIS3*, *HIS4*, *HIS5*, *HOM3*, *HOM6*, *ILV1*, *ILV2*, *ILV5*, *INO1*,
20 *INO2*, *INO4*, *lacZ*, *LEU1*, *LEU2*, *LEU4*, *luciferase*, *LYS2*, *MAL*, *MEL*, *MET2*, *MET3*, *MET4*, *MET8*, *MET9*, *MET14*, *MET16*, *MET19*, *OLE1*, *PHO5*, *PRO1*, *PRO3*, *THR1*, *THR4*, *TRP1*, *TRP2*, *TRP3*, *TRP4*, *TRP5*, *URA1*, *URA2*, *URA3*, *URA4*, *URA5* and *URA10*.

25 63. The method of claim 56, wherein said quorum sensing signal molecule is produced by a second cell.

64. The method of claim 63, wherein said second cell is a prokaryote or eukaryote.
30

65. The method of claim 64, wherein said second cell is a bacterium.

66. The method of claim 56, wherein said cell is a prokaryote or eukaryote.

35 67. The method of claim 66, wherein said cell is a bacterium.

68. The method of either of claims 65 and 67, wherein said bacterium is a gram negative bacterium.

69. The method of claim 68, wherein said gram negative bacterium is *Pseudomonas aeruginosa*.

5 70. The method of claim 67, wherein said bacterium is a mutant strain of *Pseudomonas aeruginosa* in which *lasI* and *rhlI* are inactivated.

71. The method of claim 65, wherein said second cell is wild type *Pseudomonas aeruginosa*.

10

72. The method of claim 56, wherein said quorum sensing signal molecule is an autoinducer of said quorum sensing controlled gene.

73. The method of claim 72, wherein said autoinducer is a homoserine
15 lactone, or an analog thereof.


74. The method of claim 56, wherein said quorum sensing signal molecule induces the expression of said quorum sensing controlled gene.

U.S. Pat. No. 7,222,222

Abstract of the Disclosure

10

FIGURE 1






FIGURE 2

FIGURE 3

	10	20	30	40	50	60	70	80	
1	GACCATGGCG	CACGATCGGG	TCGATGGATG	TGTAGTCTTC	GGAGATATAT	CTGCTTTTCC	ATTCCCCTGG	GTAATTGGAC	80
81	AGGAAATGAT	ATTTAGGCGC	GGTCAGCGGG	AAAGGCGCCC	GCGCACCGAA	TGAAAAGAAT	TGGAATCCAT	AATTACCGCA	160
161	TATCTCGAGA	ACCAGGGAGA	AGAACTCTTC	CTCGGTTGTT	ATTCTTGACA	AAATCTCGAG	ATATCCCTCT	CTCTCATCAT	240
241	GCATGCCAGC	TTCACCAGTC	AACAATCGAT	CCCGGCAGGA	TTCGGTTCTT	CAGCGTTTTT	ATTCTTTTTT	GTTTCCGTCF	320
321	TATCCACTCA	TGACAATTCC	CTTTATCAAG	CCGGACCTGG	AAGTACATAT	CGCCACCGGC	CCTGTGCGGG	TGTTCCGGAC	400
401	GAGGTCGAGC	CCATTGCGCG	CCCGCCGGGC	GACGACGAAA	CGAGGCTGAA	GAGAACTTCT	CGATGAGCAC	GGACACGAGG	480
481	CGCCGCGCAG	GAAGTGCCT	GGGCTGGAGG	GGACGGCGCC	CCGGATCTTT	GCGGAAACCG	TAGAACGGCT	CTCCGATGGG	560
561	CCTCACGGCG	GTCTTTCTCA	TTGTTCTTCT	CGCCAGCTCC	ATCGTTCAGT	GGAGAGCCTC	CCGACAACAA	ACCTGGCCCC	640
641	ATGGCGGGCC	TCGTCGACGA	GGCTCCCCGG	GGACCAGAAA	TGGCAACTAC	ATACTTCCCC	CCCTATCTCT	CCGCAGATAC	720
721	CTGCCCCGAA	GGGCGAGTTG	TCCCTGCCCG	GCTGTGACAA	TTTAATTGCA	CCAGGCATTT	CATTGTCCGT	GCCGATTTTC	800
801	ACGAAGCGCA	TTCTGAGGCA	ATTAAAAAGA	GCGCTCCATT	CGACCATGGA	CAAGCTATCC	ACGCCTGACC	GAGATCGCCT	880
881	TCCGAATATA	TCGAAGCGAT	AACCGCAGCC	CATCCAGCAAG	TGCTTCAGTC	AATAAACAGG	ACGCTGGCCT	TTCTGTCCGA	960
961	TGAAAGTTCC	GCATGGCGTC	CGCCCCTAAG	GAAGAGGAGA	TAAATATGAT	TTATTACTTG	ATCGGAGTGG	CGCTATTTCAT	1040
1041	CTTCATGCTG	GAACAGTTGG	TTCCCGGCTG	GAAATTGCCC	AAGGTGAGCA	CCTGGGTGGC	CCGGGTGATC	TTCTCAACA	1120
1121	TGCTCCAGGT	GTCGATCGCG	CTGCTCGCCG	GCATCACCTG	GAACAAATGG	ATGATGGGGC	ACAGCCTGCT	GCACACCTCG	1200
1201	GATGCCCTGC	CACCACTGCT	GGCCGGCTTC	GCTGCGCTACT	TGCTCAACAC	CTTCGTCAAC	TACTGGTGGC	AGTCGCGCGT	1280
1281	CCACGCCAAC	GACACGCTCT	GGCGGCTGTT	CCACCACTTG	CACCACGCGC	CGCAACGCAT	CGAGGTATTC	ACCTCCTTCT	1360
1361	ACAAGCATCC	GACCGAGATG	GTCTTCAACT	CGCTGTGTTG	CAGCTTCGTC	GCCTACGTGG	TGATGGGCAT	CAGCATCGAG	1440
1441	GCGGGCGCCT	ACTACATCAT	GTTCCGCCG	CTCGGCAGAA	TGTTCTACCA	CTCGAACCTG	CGCACCCCGC	ACGTCTCTCG	1520
1521	CTACCTGTTT	CAGCGCCCGG	AGATGCACCG	CATCCAGCAC	CAGCGCGACC	GTACAGAGTG	CAACTACAGC	GACTTCCCGA	1600
1601	TCTGGGACAT	GTTGTTTCGGC	ACCTACGAGA	ACCCCCCGCG	CATCGACGAG	CCGCAGGGCT	TCGCCGGCGA	CAAGGAGCAG	1680
1681	CAGTTCGTCG	ACATGCTGCT	GTTCCGCGAC	GTGCACAGCC	TCCCGGGAAA	AACCCAGCCC	GCTCCCGTCC	TGGTCAAGCC	1760
1761	CGACGTCAAG	TGAAGTGGTC	CCGACCATAC	CTATGCGCAA	GTCTCAGCCG	GAACATATTG	AAATCCATCT	CGTATGGCCT	1840
1841	CCCCCTCGCC	GAAAGTGGTC	CCGACCATAC	CTATGCGCAA	CTGGAACACG	GCCTCGGCGA	ACTGAAACGC	AGGTATCTGG	1920
1921	AGCTGCGCAT	CTCCACGGCG	GCGCGCGAGC	TGCCGTTTCA	CAACTACCTG	TTCTACCTGA	TCCTCCAGTC	GCGCCACCAG	2000
2001	GAATTCGACT	TCAAGCTGCG	CCAGGGCAAC	TCGGTGGTCA	CCACATATCA	CCGATTCAAG	AGCAAGGGAC	GCATCCCGTC	2080
2081	CTGACACACC	CTGCTCCTGG	CCGATGCGGT	CAACGCCAAG	AGCGAGCTGG	AGCTCAAGCA	TCCGGACATC	CCGCAGCTCG	2160
2161	ACCGCCACGC	TCGCGACATC	GAGCGCTGGC	TGGCCCGCGG	CAACGTCATG	CCGCCACGCG	AGCGGGCCCT	GCGCGGCTCG	2240
2241	GTTGAGGCGC	TGGAGCGCGC	CGCTGGCGAA	GGCCGTCCGT	TGCACCTGGT	GAGCGCGGTA	TGCCCGGACT	ACTCGCACTC	2320
2321	CAGCGATGCC	GAGGGCAAGC	CGCGCTACAC	CTTCGAGCGA	GTCCGGCGACC	AGCCCCGCCCT	GGCCGGCGCC	AAGCTGGTCA	2400
2401	GCGCCGCGCA	GCGGCTGGCG	GAGCTGGCCA	GGCGCGCCA	GGTGGAAATC	CGCCACGCGA	TCCTCGCGCG	CGAGTTCGAG	2480
2481	TACCTATCGT	TCAACCGCAA	CCCCGCCACC	GGCGAGACCC	GCGAGGGTTT	CCTCGGCAAG	GTGAGCGGCC	AGCTCGAGCG	2560
2561	GATCGCCGGG	GCCCTGCCCT	GCCCGGCCCG	GACCTGCTCG	TTCTTCGAGA	TGTGCGGCGG	CGAGGACGGC	TGGCACCAGG	2640
2641	CCCACGGCGA	GATCGTCCAG	CGCCTGGAAC	AGGGCGACTA	CGGCCAGACC	GGGCTGGACT	ACCCGGCCCT	GGAATCGATC	2720
2721	TTCCTGTGCG	GCCTGCCGCT	CTACGAGAAA	TGTTTCGCCA	GCCAGTCCGG	CGAGCAGATC	TGGGCCAGCT	TCGTCTCCCA	2800
2801	GGCCGCCGAG	TACGCATTGA	TGGGAAAAC	CTTCGGCGAG	CGCTTCGACA	ACTTCGTCTG	GCTGGCCGTC	GATCACTACC	2880
2881	GGATGGAGCC	GTTCTACTCG	TTCTTCGCGA	CCGTCCCGAC	GCTCTACATC	CGAACCAGACT	ACCTGTAACG	AGGGAGCGCC	2960
2961	TGCGCCATGC	AAGATGAAC	GTTCAAGACC	CGATACTCCA	AGTACGGATA	CGGCATCGAC	GTGCGCCGTA	CCTACAAGGA	3040
3041	CCTGCCCTGC	CAGCGTTTCT	GGACCTGGGT	CACCGGCAAG	TCGCTGAACG	ACCGCCCGCC	GCGACGGCCG	AAAGACAGCC	3120
3121	TGCTCAAGCC	CTGGCAGCTC	TACCTGCACA	TCAGTTGGGG	CTACGCGGTG	TTCTTCCTCG	CGGTGATCTA	CGGCCAGCAA	3200
3201	CTGCTCGCCT	CGCAGCAGCC	ATTGTGGCTG	AAGTGCTTGC	TGGCGCGGTT	GATCATGTGC	CTGGTGGTCA	ACCGCCAGCG	3280
3281	TGGCTTCTCT	CATACCTTCC	ACTACACCAC	CCATGGCGCC	AGCCTGGAGA	ACAAGGCGCT	GGCCCGCTTC	ACCTGCAAGT	3360
3361	GGATCCTGTC	GATCCCGATC	CTGCACACCC	CGCGCGACGA	GTACGTGAAG	CTGCACGTGA	ACGAACACCA	CAGCGTGC	3440
3441	ACCTTCAATA	CCGAGCACGA	CGTCGACCTG	GTCTTCATGA	AACAGCACGG	CTTCTACAAG	GGCATGTCCG	AGAGCGCCTT	3520
3521	CTGGACCCCG	CTGGTGCTCG	CGCCCTTCCA	TCCGGCGCGG	ATCCTCGAGC	ACCTGAAGTT	CCGTTTCGAC	GTGAGCTTCG	3600
3601	TCTCCGCCCC	GCGCCACGAG	CGCGTCAGCC	GGGCGCTCTA	CTGGGCTGCG	CTGCTCGGCC	TGGTGACGCG	CAGCGGCTAC	3680
3681	CTGGAGGCGT	TCGCGCTGTT	CTACCTGTTT	CCGATCTTCA	TCCTCACCCA	GTAAGTCTCG	TGGATCCAGC	ACGTCTCCGA	3760
3761	GCACCTCTGG	TTGCGCCGCA	ACGAGCACGG	CCTGCCGCGC	TTCTTGCACT	ACGGCTCGCT	GAGCTGGGGA	CGCTTCTCTG	3840
3841	GCCGCCCTTA	CCCGGCCGAC	AAGCAGGGCC	TGGCCTTCGC	CCTGGCGTTC	GTTTCGCTGGA	GCCTGGGCGT	GCTGCTGATC	3920
3921	GACATCCCGC	TGCGGGTGTT	CTCCTTCATG	CAGGACCTGC	CCAGCCACGA	CTTCCACCAT	CGCAAGCCGG	GAGTGAACCT	4000
	10	20	30	40	50	60	70	80	

CCCTCCAGGT





FIGURE 4

FIGURE 5






FIGURE 6

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted October 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

qec102	A	C	C	T	G	C	C	C	T	G	A	A	G	G	G	C	A	G	G	T
qec104	A	A	C	T	A	C	C	A	G	T	T	C	T	G	G	A	A	G	G	T
qec117	C	A	C	T	G	C	C	A	G	A	T	C	T	G	G	C	A	G	T	T
qec126	A	C	C	T	A	A	C	A	G	A	T	T	T	G	T	A	A	G	T	T
qec128	A	C	C	T	A	C	C	A	G	A	T	T	T	G	G	C	A	G	G	G
qec131	A	C	C	T	A	C	C	A	G	A	T	C	T	T	G	T	A	G	T	T
qec132A & B	C	C	C	T	A	G	G	C	G	G	C	C	T	T	G	G	A	G	C	C
rA/AB	T	C	C	T	G	T	G	A	A	A	T	C	T	G	G	C	A	G	T	T
rA/I	C	C	C	T	A	C	C	A	G	A	T	C	T	G	G	C	A	G	G	T
laxB	A	C	C	T	G	C	C	A	G	T	T	C	T	G	G	C	A	G	G	T

FIGURE 7

FIGURE 8






FIGURE 10A





FIGURE 10B



09063730 : 090400

FIGURE 11





FIGURE 12



Customer Number: 000959

Attorney's
Docket
Number UIZ-038

Declaration, Petition and Power of Attorney for Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

QUORUM SENSING SIGNALING IN BACTERIA
the specification of which

(check one)

X is attached hereto.

 was filed on _____ as

Application Serial No. _____

and was amended on _____
(if applicable)

I do not know and do not believe that the subject matter of this application was known or used by others in the United States or patented or described in a printed publication in any country before my invention thereof, or patented or described in a printed publication in any country or in public use or on sale in the United States more than one year prior to the date of this application, or first patented or caused to be patented or made the subject of an inventor's certificate by me or my legal representatives or assigns in a country foreign to the United States prior to the date of this application on an application filed more than twelve months (six months if this application is for a design) before the filing of this application; and I acknowledge my duty to disclose information of which I am aware which is material to the examination of this application, that no application for patent or inventor's certificate on the subject matter of this application has been filed by me or my representatives or assigns in any country foreign to the United States, except those identified below, and that I have reviewed and understand the contents of the specification, including the claims as amended by any amendment referred to herein.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

CLAIM OF BENEFIT OF EARLIER FOREIGN APPLICATION(S)

I hereby claim priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below, and have also identified below any foreign application(s) for patent or inventor's certificate filed by me on the same subject matter having a filing date before that of the application(s) from which priority is claimed.

Check one:

☒ no such applications have been filed.

☐ such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No

ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

CLAIM FOR BENEFIT OF U.S. PROVISIONAL APPLICATION(S)

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

60/153,022
(Application Serial No.)

September 3, 1999
(Filing Date)

(Application Serial No.)

(Filing Date)

CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S)

I hereby claim the benefit under Title 35, United States Code, §120 of any earlier United States application(s) or PCT international application(s) designating the United States listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the earlier application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date(s) of the earlier application(s) and the national or PCT international filing date of this application. As to subject matter of this application which is common to my earlier application(s), if any, described below, I do not know and do not believe that the same was known or used by others in the United States or patented or described in a printed publication in any country before my invention thereof, or patented or described in a printed publication in any country or in public use or on sale in the United States more than one year prior to the date(s) of said earlier application(s), or first patented or caused to be patented or made the subject of an inventor's certificate by me or my legal representatives or assigns in a country foreign to the United States prior to the date(s) of said earlier application(s) on an application filed more than twelve months (six months if this application is for a design) before the filing of said earlier application(s); and I acknowledge that no application for patent or inventor's certificate on said subject matter has been filed by me or my representatives or assigns in any country foreign to the United States except those identified herein.

_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented,pending,aband.)
_____ (Application Serial No.)	_____ (Filing Date)	_____ (Status) (patented,pending,aband.)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

James E. Cockfield	Reg. No. 19,162	Nicholas P. Triano III	Reg. No. 36,397
Thomas V. Smurzynski	Reg. No. 24,798	Peter C. Lauro	Reg. No. 32,360
Ralph A. Loren	Reg. No. 29,325	DeAnn F. Smith	Reg. No. 36,683
Giulio A. DeConti, Jr.	Reg. No. 31,503	William D. DeVaul	Reg. No. 42,483
Ann Lamport Hammitte	Reg. No. 34,858	David J. Ridders	Reg. No. 43,882
Elizabeth A. Hanley	Reg. No. 33,505	Chi Suk Kim	Reg. No. 42,728
Amy E. Mandragouras	Reg. No. 36,207	Maria Laccotripe Zacharakis	Limited Recognition
Anthony A. Laurentano	Reg. No. 38,220		Under 37 C.F.R. § 10.9(b)
Jane E. Remillard	Reg. No. 38,872	Debra J. Milasincic	Reg. No. P46,931
Jeremiah Lynch	Reg. No. 17,425	David R. Burns	Reg. No. P46,590
Kevin J. Canning	Reg. No. 35,470	Sean D. Detweiler	Reg. No. 42,482
Jeanne M. DiGiorgio	Reg. No. 41,710	Peter S. Stecher	Reg. No. P47,259
Megan E. Williams	Reg. No. 43,270		

Send Correspondence to Peter C. Lauro, Esq. at **Customer Number: 000959** whose address is:

Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109

Direct Telephone Calls to: (name and telephone number)

Peter C. Lauro, Esq., (617) 227-7400

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Marvin Whiteley	
Inventor's signature	Date
Residence 991 22 nd Avenue, Coralville, IA 52241	
Citizenship U.S.A.	
Post Office Address (if different)	

Kimberly M. Lee

Date _____

1162 Briar Drive, Iowa City, IA 52240

U.S.A.

Post Office Address (if different)

E. Peter Greenberg

Date

4020 Stewart Road, Iowa City, IA 52240

U.S.A.

Post Office Address (if different)

Ute Muh

Date

410 North 7th Avenue, Iowa City, IA 52245

Germany

Post Office Address (if different)

SEQUENCE LISTING

<110> Whiteley, Marvin
 Lee, Kimberly
 Greenberg, E. P.
 Muh, Ute

<120> Quorum Sensing Controlled Genes in Pseudomonas
 Aeruginosa

<130> UIZ-038

<140>

<141>

<150> 60/153,022

<151> 1999-09-03

<160> 39

<170> PatentIn Ver. 2.0

<210> 1

<211> 1218

<212> DNA

<213> Pseudomonas aeruginosa

<400> 1

```

atggacgatac tattgcaacg cgtacggcgc tgcgaagcgc tgcagcaacc cgaatggggc 60
gatocgtcgc gcctgcgcga cgtgcaggcg tacctgcgcg gcagtccggc gctgatccgc 120
gocggcgaca tcctggccct gcgcgcgacc ctggcgcggg tcgcccgcgg cgaggcgtg 180
gtggtacagt gcggcgactg cgcgcaggac atggacgacc accatgccga gaacgtggcg 240
cgcaaggccg ccgtgctgga actgctggcc ggcgcctgc gcctggccgg ccggcgcccg 300
gtgatccgcg tcgggcgcgt cgcggggcag tacgccaagc cgcgttcaa gccgcacgag 360
caggctggcg agcagaccct gccggtctat cgcggcgaca tgggtcaacg ccgcgaggcc 420
catgccgaac agcgcggggc cgatccgcag cggatcctca agggctatgc ggcgcgcgcg 480
aacatcatgc gccacctggg ctgggacgcc gcgtccgggc aggaggcgaa tgcctcgccg 540
gtctggacca gccacgagat gctgctgctc gactacgagc tgtcgatgct gcgcgaggac 600
gagcagcgcc ggggtctatct cggttcgacc cactggccgt ggatcggcga gcgcacccgc 660
caggctcgacg gcgcccattg ggcgctgctg gccgaggtgc tcaaccgggt ggcctgcaag 720
gtcgggtccgg agatcgcccg cgaccagttg ctggcgctct gcgagcgccg cgatccgcgc 780
cgcgagccgg gacgcctgac gctgatcgcg cggatggcg cgcagaaggt cggcgagcgc 840
ctgcgcgcgc tgggtggaggc ggtgcgcgcg gccgggcacc cggatgatctg gctgagcgag 900
ccgatgcacg gcaacacat cgtcgcgcgc tgcggcaaca agaccgcct ggtgcgcagc 960
atcgccgagg aggtggcggc gttccgcctg ggatcctcta gccgaggcgg cgtgctcaac 1020
ggactgcacc tggaaaccac cccggacgac gtcaccgagt gcgtcgccga ttccagcggc 1080
ctgcaccagg tcagccggca ctacaccagc ctctgcgata cgcggctgaa ccctggcag 1140
gcgtcagcg cggatgatggc ctggtccggc gcagaagcga tccagagcgc aaccttcccc 1200
ctggagaccg tggcatga                                     1218

```

<210> 2

<211> 1782

<212> DNA

<213> Pseudomonas aeruginosa

<400> 2

```

atggatgatg gggcacagcc tgctgcacac ctcgatgccc ctgccaccac tgctggccgg 60
cttcgccgcc tacttcgtca acaccttctg cactacttgg tggcatcgcg cgcgccacgc 120
caacgacacg ctctggcggc tgttccacca gttgcaccac gcgccgaac gcacgaggt 180
attcacctcc ttctacaagc atccgaccga gatggtcttc aactcgctgc tgggcagcct 240

```

U.S. PAT. & TM. OFFICE

cgtcgcctac gtggtgatgg gcatcagcat cgaggccggc gcctactaca tcatgttcgc 300
 cgcgctcggc gagatgttct accactcgaa cctgcgccacc ccgcacgtcc tcggctacct 360
 gttccagcgc ccggagatgc accgcatcca ccaccagcgc gaccgtcacg agtgcaacta 420
 cagcgacttc ccgatctggg acatgttggt cggcacctac gagaaccccc gccgcatcga 480
 cgagccgcag ggcttcgccg gcgacaagga gcagcagttc gtcgacatgc tgctgttcgg 540
 cgacgtgcac agcctccccg gaaaaaccca gcccgtctcc gtccctggta agcccagct 600
 cagggtgaacg ccattgattcc agacatcgat tcccgctctca gccggaacat attgaaatcc 660
 atctcgtatg goctccccct cgccgaagtg gtccccgacc atacctatgc gcaactggaa 720
 acgcgcctcg gcgaactgaa acgcaggtat ctggagctgc gcatctccca cggcgcgcgc 780
 gagctgcggt tcagcaacta cctgttctac ctgatcctcc agtcgcgcga ccaggaattc 840
 gacttcaagc tgcgccaggg caactcgggt gtcaccaaca tccaccgatt caagagcaag 900
 ggacgcaccc cgtccctgac caccctgtct ctggccgatg cgggtcaacgc caagagcgag 960
 ctggagctca agcatccgga catcccgag ctcgaccgcc acgctcgcga catcgagcgc 1020
 tggttgccg ccggcaacgt catgccgccc agcgagcggg ccctgcgcgc cctggttag 1080
 gcgctggagc gcgcgcgtgg cgaaggcgt ccgttgccacc tggtagagcg ggtatgccg 1140
 gactactcgc actccagcga tgcgcagggg aagccgcgt acaccttcga gcgagtcggc 1200
 gaccagcccg gcttgccgg cgccaagctg gtcagcgcgc gccaggcggg gccggagctg 1260
 gccagggcgc gccagggtga aatccgccac gcgatcctcg gcggcgagtt cgagtacct 1320
 tcgttcaacc gcaacccgc caccggcgag accgcgagg gtttcctcgc caaggtcgag 1380
 cgccagctcg agcggatcgc cggggccctg ccctgcccg ccgcgacctg ctctgttctc 1440
 gagatgtcgc gcggcgagga cggctggcac caggcccacg gcgagatcgt ccagcgcctg 1500
 gaacagggcg actacggcca gaccgggctg gactaccgg ccctggaatc gatcttctg 1560
 tcgcgcctgc cgtctacga gaaatgggt gccagccagt cgcgcgagca gatctgggc 1620
 agcttcgtct ccaggccgc cgagtacga ttgatggaa aactcttcg cgagcgcttc 1680
 gacaacttcg tcgtgctggc cgtcgatcac taccggatg agccgttcta ctctgttctc 1740
 gcgaccgtcc cgacgctcta catccgaacc gactacctg aa 1782

<210> 3

<211> 693

<212> DNA

<213> Pseudomonas aeruginosa

<400> 3

atgcgcgcaa ccagccccac accaaccaac ccgcatccca ggctaccgcc tgatcacaca 60
 ggaaacccca tgaatactca gattgccag atcaccaga gcctggcagc caacggctgc 120
 gcctatatca cccccagcga cgcgctctac gacgagcagg actgggaact gatgaaccag 180
 gtccctggca actcgacct gccgtgggag aagatcctga tcggcgacgc cgacgaggag 240
 aacgaacctt acgtggcccg tttcatgacc gaccgcgacc gtcccacggt ggtcaaccat 300
 gcgctgtcgg agctgatcat cccgcgcgtc tgcaacgaca acgtgatgag cctgttccgc 360
 aagctgatgg gcgacgacgc cttctacgtc cggcgcatgc aggtgaaccg gatgaaggcc 420
 ggctcgttca tcggccggca cctggatacc gacagcaacc cggactacca gtactccatc 480
 gtccctgcagc tcggcaccta cttctccggc ggccagttcg tggctacga ccgcgacggc 540
 aacctgcgca acgacatcaa gccggagccg cgtcgggtga tcatcagcga ctgtagctat 600
 cccacagagg tccagcaggt gaccgccggc gacgcgctct cgctggtgtt cttcgtcagt 660
 cgccatgcgg accggaaccg gcgggtctat tga 693

<210> 4

<211> 411

<212> DNA

<213> Pseudomonas aeruginosa

<400> 4

gtgacggact tcgaatcctc gcgtcgcgt ccgtccaccg gattgtccgg cgcgctgcgg 60
 cgccgcgct ccagcgcagc gccactgcca gatctggcag ttgtcgtgg cggggacgga 120
 ggcttttagta gccgcacttt tttccagggc cgggcagtgg gaccgcaatg catggacgac 180
 atcgagacca gactgaggaa actggtagcc gcccggttcg gcgtggagga atgcgacatc 240
 cggttgga gacacttccg taacgacttc ggtgccaggt cgctcgaggt agtcgaactg 300
 gtcattggccc tggaagcgga gttcggcgtc gagatcgcc atgacgatgc ggaacggatc 360
 gagaccgtgc gccaggccat cgactatctc gaggaagccg tgccgacctg a 411

2025-06-10 10:00:00

<210> 5
 <211> 588
 <212> DNA
 <213> *Pseudomonas aeruginosa*

<400> 5
 atggccgtcc gggctcagga agtagaacga atgcacctcg ctgcggttct gcttccattc 60
 gcgcacgcc tgcgcgcga gctgcgcggc gaagcgggcg aaatccgcgg cggcgatgcc 120
 gaaggcgtag tgcgtgtagt ccgcggccgg cccgcggtac tgcggctccc gggacaggca 180
 cagccacagc gaaccaggtt cgagataggg gccctggtcc cagcgcgctt ccaggcgaaa 240
 gccgagaaga tcgcggtaga aggcgatgct ggccggcagg tcggcgaccg ccagggtcag 300
 gtgattgaga ccggttaagca tgggggctcc ttgcaagatg tggcgggagg tcgattcagg 360
 cacgtcccag ccagtcgccc cggatcattt ccatcagttg gcgcaagccg ggttgcggtc 420
 ggcgtcggtc cggatagtag aggcagaacg gcgcgcccac cgaggtccag tccggcaata 480
 ccagttgcag ccggccgcta cgcagctcct cggcgattcc cacctccagg cagtaggcca 540
 ggccgacacc gtccagggcc gcggcaaccg ccgtattgct ttcgttga 588

<210> 6
 <211> 1020
 <212> DNA
 <213> *Pseudomonas aeruginosa*

<400> 6
 atgaacgaa ccgcgcgcga taccctcgcc gtatcgcccc cgccctcgcg caacctctgc 60
 gacggccacg gccggctcga tccccgggcg gtccggtggt ccgcccggcc gcgggtgctc 120
 tgcacatcc ccggccactt ccggccggcg aagcgtgga accactggtg catcgtcagc 180
 cccggtgga tgcctctgct gaccatcgcc gacctcgact acctgacctt cggcgccgcc 240
 tatttctctg acctggacag cggccaggcg gttagcgaca cgcagatccg cttcttcggc 300
 ctccggtgcc agttgcccga cgcgcgcgag gccagccatg ccttcgagca tccccgcctg 360
 caattgcgct tcgacgaaca gcccgggcgc ctgcgcgtca ccggccaggc cccggacctc 420
 ggtggcctgc cgctggagct ggcgctgga gtgcgacgac cgctgcacct ggagtcggtg 480
 aacctggtgg tgcgatggg cgaacacacc ttccatgcct gcagccgcca gctcggcctg 540
 ccgatcagcg gctgctgca gctcgccgc cgacgctacg actgccaggc gggccagagc 600
 ttccgcgcgc tggacttcgg ccgcggtgct tggccgctgc atacctactg gaccgcgcgc 660
 gccttcgccc ccccgggcgg catcgccggc aacttcggca ccggctggac cgaagccagc 720
 gacctgcgcg agaacgccct gtggttcggc ggcaagctca gccgcgtgct cgacgacgtg 780
 cacatccgcg agcctcgcga cccgctggcc gaatggcgcc tggacagcgc ctgcggtcgc 840
 gtccagctgc tcttcgctcc cgaacagctg caccaggcgc ggcccagcgt cggcctgttc 900
 tatgccata cccgccagtg gttcgccgct ttcaacggca ccctgcgcca cgacgacggc 960
 gactgcgtgc cgggtggacgg cgccctcggc tggatcggtt cgaccgcgc gcgctggtga 1020

<210> 7
 <211> 1170
 <212> DNA
 <213> *Pseudomonas aeruginosa*

<400> 7
 atgaaacact actcagccac cctggcactc ctgccactca ccctcgccct gttcctgccc 60
 caggcagccc atgcccacgg ctccatggaa acgcgcgcca gtccgggtcta cggtgcttc 120
 ctogaaggtc cggagaatcc caagtcggcc gcttgcaagg ccgcgcgtcg ccgcggcgcc 180
 acccaggcac tgtacgactg gaatggcgct aaccagggca acgccaacgg caaccaccag 240
 gcggtgggtc ccgacggcca gctctgcggc gccggcaagg cactgttcaa gggcctgaac 300
 ctggctcgca gcgactggcc cagcactgcc atcgcgccgg acgccagcgg caacttccag 360
 ttctgtctaca aggcacagcg gccgcacgag acccgctact tcgacttcta catcaccaag 420
 gacggctata accccagaaa gccgctggcc tggagcgacc tggaaaccgc gccgttctgc 480
 tcgatcacca gcgtcaagct ggagaacggc acctaccgga tgaactgcc gctgccccag 540
 ggcaagaccg gcaagcatgt gatctataac gtctggcagc gctcggacag cccggaagcc 600
 ttctacgctt gcctcgacgt gagcttcagc ggcgccgtcg ccaacccttg gcaagcgtg 660
 ggcaacctgc gcgcgcagca ggacctgcca gccggtgcta ccgtcaccct gcgtctgttc 720
 gatgcccagg gccgcgacgc ccagcgtcac agcctgacct tggcccaggg cgccaacggt 780

2025 RELEASE UNDER E.O. 14176

gccaaagcaat ggccgctggc gctggcgagc aaggtcaacc aggactccac cctgggtcaac 840
 atcgggcgtgc tggatgccta cggggcggtc agcccggtgg ccagctcgca ggacaaccag 900
 gtctacgtgc gccaggccgg ctaccgcttc caggtcgaca tcgaactgcc ggtcgagggc 960
 ggcgggcgagc aaccggggcg cgacggcaag gtcgacttcg actatccgca aggcctgcag 1020
 caatacgacg ccgggaccgt agtgcgcggt gccgatggca agcgctacca gtgcaagccc 1080
 taccogaact ccggctgggt caagggtctg gacctctact acgccccggg caagggcag 1140
 gectggcagg acgcctggac cctgctgtaa 1170

<210> 8

<211> 210

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 8

atgttgaaag tggcgatcgt cctgctactg ctggctaccc tgggtgagcct gttcagcggc 60
 ctgttcttcc tggtaagga ccagggccat ggttcccgcg tggtaattc gctgaccgtc 120
 cgctgtgtgc tcgcccggc gaccctggtg ctggctgcct ggggcttcta cagcggcag 180
 ctgaacagcc acgcgcctg gcatttctga 210

<210> 9

<211> 1872

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 9

atgagtttcc cgataaacat caattatagg agtttcccta tgtcggtct cgcggttgg 60
 gtggattaca cgcgcaagct cgacgacgaa tttccggcga tcttcgccat gaccgatacg 120
 ctgccttgc gggggccgga tgccgagggc atctggaagc accgcaacgc cctgctgggt 180
 caccggcggc tggcggtcat cgacctcagc ggcgcggtgc agccgatgtc ctatcgcttt 240
 cccacgggcc aggaggtcac cctcgtctac accggcgagg tgtacaacca cgatgccctg 300
 cgcgagcggt tgcgcggggc cggacatgag ttccgcaccc gcagcgatac cgaggtggtc 360
 ctgcacgcct atctgcaatg gggcgagcgt tgttgcgagt acctgaccgg gatgttcgcc 420
 ttgcgcgtct togatggccg cgacggccac ctgctgctgg tgcgcgaccg cctgggcatc 480
 aagcgctgt attacgcgcg gcaccgcgag ggaactgctgt tcggctcgga gatcaagtcc 540
 atcctggcgc atccggaatt cgccgccagg ctgcagcgcg tcggcctggt cgacctcctg 600
 acgtgtccc ggggcacttc gcagacgcg ttccgcgagg tccaggaact gctgcccggc 660
 caoctgctgt cctggcgctc caattcccag gcgaagtgc gccgctactg ggaggtgcgc 720
 cgccaggagc atgcgacga cctgcagagc accgtgcagc gcaccccgca actggtcacc 780
 cgcgccctgg gggcgcaatt gcacgccgac gttccgggtg gttcgtgct atcggtggg 840
 ctcgattcga ccgccctgac cggcatcgcc cagcgcatcg cgaaggcgga gcacggcggc 900
 gacatcaatt cgttctcggt ggacttcgtc ggccaggccg agcagttccg cagcgacgac 960
 ctgcgtcccg accaggacca gccgttcgcc ctgctggccg cgcagtaacat cggcagccgt 1020
 catcgaccg tgctcatcga caatgccgaa ctggtctgcg aacgagcgcg cgaagaggta 1080
 ttccgggcca aggacgtacc ttacaccttc ggcgacatgg atacctcgt gcacctgatg 1140
 ttccggcgaga tccgcgggca ttccacggtg gccatctccg gtgaaggcgc cgacgagctg 1200
 ttccggtggct acggctgggt ccgcgatccg caggcggtgg ctgcggcgcg cttcccctgg 1260
 gcctccaggg tgcgcctgcc ggccggcttc atcgacgccg gtttcaaccg ccgctgcgat 1320
 ctctccagt accagcaggc cagctacgac gatgggctgc gccaggctga acacctggcc 1380
 ggcgacagcc cggaggagcg gcggatgcgc gacttcagcc acctgcatct gaagcgctgg 1440
 atggtgctgc tgcctgaacg caaggatcgc ctgagcatgt gcaacggcct ggaggtgcgg 1500
 gtgcctaca ccgaccatga gctgggtggag tacgtctaca acgtgccctg gtcgatcaag 1560
 agccgggacg gcgaggagaa gtgctgctc aagcgggct cgcgcgacta tgtcccgaa 1620
 gccgtgctca agcgcgcgaa gagcccttat ccgacttctg ccaacctcgg ctacgagcgt 1680
 ttctgcgcg ggagcgtgcg gcgcctgctg gaggacgcgg tgaaccgggt gttcggcatc 1740
 gtttcgcgag agttcctggc ccgcgaactg gagcatccgg aggggtactt caacaccag 1800
 gtgagccgcc acaacctgga gaccgcactg gcgctggaag gctggctcag gttgtacggg 1860
 ctctccgct ga 1872

<210> 10

<211> 756

<212> DNA

<213> Pseudomonas aeruginosa

<400> 10

```

atgcagaaac agcgggtggc cgaccaggtc gcagagcgta tcgagcgggt gatcgtcgac 60
ggcgtgctca aggtcggcca ggcaactgcc tccgagcggc gcctgggtggc caagctcggc 120
tgctcgcgct cggcoctgcg cgagggcctg cgggcgctgc gcgggcgcgg catcatcgac 180
accgagcatg gccgtgggtc gttcgtcgcc gacctcgacc gcaacgccga cgtcagcccc 240
ctgatgcacc tgttcggctc ccagccgcgc acctctacg acctgctcga agtccgcgcc 300
ctgctggagg gcgaggcggc ccgcctggca gcgctacgcg gcaccgaggc agacttcgtc 360
ctgctcgcgc ggcgctacga agagatgctc gccagccacg aggaaaccca gccgatcgat 420
ccccgcgagc acgcccgcgc cgaccacgcg ttccaccggg cgatcagcga ggcacgcgac 480
aatccgggtg tgggtcatac cctgcaatcg ctcaacgaac tgctgctgag cacggtgttc 540
gcctcagtga acaacctcta ccaccgaccg ccgcagaaac gccagatcga ccgccagcac 600
gcgcgcctct acgcggccct ccgcgagcgc cagccggacc aggcgcaacg ggcggcgcgc 660
gaacatatcc acagcatccg cgacaacctg cgggagatcg agcaggaaga acagcgctcg 720
gtccgcgcca ccctgcgcct gaacggctgg ggtga 756

```

<210> 11

<211> 822

<212> DNA

<213> Pseudomonas aeruginosa

<400> 11

```

atgaaccatg tcatcaccoc ccacagcaag ctgctcggcg tcatcgagcc ggtcctcaac 60
gacatgcccg ccggaacctt gcgcaacgca ctgttcgggg ccttctggga cgagacggcg 120
tcgttgctgg acatcgagga cgccttcgcc cgggtcaccg cccggcgcca ggcggtcgag 180
ccgctgcgca agttcttcgc cagttggctg aagaccaaca actcggcggc cagcgtttcc 240
ggactggcca atcgcccttac cctgctggcc cgttcggaac agggttcggc agcggcagac 300
cagctctatc gagcctgcgg cagcctgcaa cggatcacgc acgaagacct cggcgccctc 360
ggcaacaccg tgcattgcga tctttcttac acctggcca ccacccttg cggcgacgac 420
cgctggctgc tgcgcgagaa ctgcctgcct tcggcgagc cgttcaagga ctggaccgac 480
cgccagcgcc tgtgcgagcg cgacctgat caggactgc tgaccacgct ggtacacgag 540
gtctatatcc acggcgaggt ggagtacatc caccgctgtt acaaggaatg gttcagccgc 600
gacatgggcg taccgcgcga acgcgcgcgc gccaccgtgg cctgggtaac ggtgcacacc 660
ggcggcaccg agagcaatca cttcgcccac gccacggcgg cggatgaacgc ctctcgtcag 720
gogatggaaa tcgaggtgaa cgaagaagcc gcgcgcaacc tggtcgggct ctacctgcgc 780
aacaaggcgc aggtcatgcg tgactgcgcg gcgctgttct ga 822

```

<210> 12

<211> 1368

<212> DNA

<213> Pseudomonas aeruginosa

<400> 12

```

atgtcctccc gccaatcgtc ccgcaacgct tccaccccgat atctgaccaa ggccttcag 60
gcaacggcca tcgtcgtggg gagctacttc ctctactgga cctaccagct ctaccagtac 120
ggcgagattc ccatcagcaa gaaggacgtg atgctgcgcc aagccatcct cgcgcgcttt 180
ccggcggaact acgaggtgga gatcaagggc gccgacctgc tcggcttcgg cgagaaattc 240
ctggctgcct acggcaatcg gcgcttcgtc ggcaaggcct tcgccatgga cgaccaggtc 300
atcgagcgcc tggagcgga ccagggacgc accaacctgc cgctggtgaa ggtgttctac 360
atcgccgaac ccggcctcct ctctcgtcgt ctcaacctct ccccgttcct ggatatccag 420
aagaacatgg tcgagctgag cctccgggaa taccggaaga tccagttggg ccccttcgat 480
ccggacgcga agcggaacc gcgcgagcag ttcgaaaccg attatgcctt ccccgagctg 540
ttcagcctca gccaaactgga agtcgcgcac tacgacggcg acggccgcga cgaactgcgc 600
ctgggctacc tgtcctacgc cggcggttcg ggagggacgc gctggctcgg gatctacgac 660
ctgaaggacg gcgcgctgac cgccattcc ggctatccgg aaatgctcga catcgacgtc 720
gcccggttca tccaggcggt caacctgtac gccggcctcg acggcacctt gccgcgcgac 780
cagcgtcagc tggagagcgt ggtcggcgga ggcagcgagc gcttcgccct gaccgcgcgc 840
gagcgccagc cactggctgc cgaccgcgc cagcgggacg actacgccag ggtcctgatg 900

```

2025 RELEASE UNDER E.O. 14176

```
<210> 13
<211> 1209
<212> DNA
<213> Pseudomonas aeruginosa
```

```
<210> 14
<211> 3090
<212> DNA
<213> Pseudomonas aeruginosa
```

<400>	14						
atgaccttta	ccgacctggt	cgtccgccgg	cgggtgctgg	cgctggtggt	cagcacgctg	60	
atcctgctgc	tgggcctggt	ctccctgggc	aaagtgcga	tccgccagta	cccgtgctg	120	
gaaagctcga	ccatcacctg	caccaccgag	taccccggcg	cctccgccga	tctcatgcaa	180	
ggcttcgtca	cccagccgat	cgcccaggcg	gtgtgctcgg	tggagggcat	cgactacctt	240	
tcctcgacct	cggtgcaggg	gcgtagcgtg	gtgaccatcc	gcatgtctgt	caaccgcgat	300	
tcgacccagg	cgatgacoga	gaccatggcc	aaggctcaact	cggtgcgcta	caagctgcc	360	
gagcgtcctt	acgactcggg	gatcgaacgc	tcttcgggcg	agaccaccgc	ggtagcctac	420	
gtcggtcttt	ccagcaagac	cctgccgata	ccggcgttga	ccgactacct	gtcgcggggtg	480	
gtcgagccga	tgtttctctt	catcgacggc	gtggccaagg	tccagacctt	tggcgggccag	540	
cgcttgccca	tgcgcctctg	gctcgacggc	gaccgcctcg	ccgggcgcgg	cctgaccgcc	600	
tccgacgtgg	ccgaggcgat	ccgccgcaac	aactaccagg	cggcgcgcgg	gatggtgaag	660	
gggcagtagc	tgctgtccaa	cgtgcgggtc	aacaccgacc	tgaccaacgt	cgacgacttc	720	
cgcgagatgg	tcatccgcaa	cgatggcaac	ggcctggtgc	gcctgcgcga	cgtcgggtacc	780	
gtcgaactgg	gcgccgcggc	caccagagac	agcgcactga	tggagcgcca	ccggcgcggtg	840	
caacctgggtg	tgttcccgac	gcccaccggc	aaccgcctgg	tgatcgtcga	cggcataccg	900	
aagctgtctc	cggagatcca	gaagaccttg	cgcgcggatg	tccgcgtcga	cctgcctac	960	
gagacttcgc	gcttcattcca	ggcctccatc	gacgaggtgg	tgcggaccct	ggtggaagcg	1020	

[illegible]

[illegible][illegible][illegible]

agcctggagc gtcggcgaa catgctgcag ggcgtctacc agcgcggcct gggcaacgac 1140
 gtcaccgcat acgcgggcgg ggccttcacc ggcagctaca tgtccgggtt gatgggcgcg 1200
 gcgctgaaca cgcgggtggg cggattctcc ggtgacgtga cgctggcgcg taccgaggtt 1260
 cccggcgacg accgccttag cggctccagc taccgtctcg cctacagcaa gaacctgccg 1320
 aacaccggca ccaacttttc gctgctcgcc tatcgttact ccaccgggtg ctatctcggc 1380
 ctgcgcgacg cggccttcat gcaggaccgg gtagagcgag gcgagccgct ggagtcgttc 1440
 tcgcgcttgc gcaatcgtct cgacgccaac atcagccagc aactgggcaa cggcggaac 1500
 ctttacctga acggctcctc gcagcgctac tggagcgcg gcgggcgggc ggtcaacttc 1560
 tccgtcggct acagcaacca gtggcgcgac gtcagttact ccatttcgcg gcaacgcctg 1620
 cgcagccagt acgaaggctt ttccagcggg gacaggcgcg gcgagaccag cacgctgttc 1680
 agcctgaacc tgtccattcc gctcggcggc gctggacgcg ggtcgccgac cctgagcagc 1740
 tacctgaccc gogacagcaa cagcggaacc cagctacca gcgggggttc cggcatgctg 1800
 ggcaagcggt gcgaggcctc ctactcgctg tcggcctccc atgaccgcca cagccggcag 1860
 acctcgaaga gcgccagcct cgactatcga ctgccgcagg tcgaactcgg ctccagcctc 1920
 tcgcagggac cgggctatcg gcagttgtcg gtcaaggcgg cggggggcct ggtcgcgcac 1980
 agcggcgagg tcaccgcggc acaaaccctg ggcgagacga tcggcctggt ccacgcgcca 2040
 aatgccaggg gcgoggtgc cggctactcg ggaagccgga tcgaccgcca cggctatgcg 2100
 gtgattccca acotgctgcc ctaccagttg aacagcgctg acctcgacc caacggcatg 2160
 gccgacgaga tcgaactgag gtccagttcg cgcaacgtgg cggccaccgc cggagcgggtg 2220
 gtgcgcctcg actatccgac gcgggtggca aggccttgc tgggtgtag ccgatgccc 2280
 agcggcgagc cctgcgctt cgccgcggaa gtgctcgatg cccacagcgg gcagtcgggtg 2340
 ggcgcgctcg gccagggcag ccgcctggtg ctgcgggtcg agcaggatcg cggctcggtt 2400
 cgggtgcgct ggggcaacga gccgcagcag cagtgcctgg tcgactatgc gttgggcccg 2460
 cgcgagacga cgctcccgct cctgcaactg gcatgtcgcc cggcgtcggc cgccgaccgg 2520
 gagcgcacgc tgtga 2535

<210> 16

<211> 2976

<212> DNA

<213> Pseudomonas aeruginosa

<400> 16

atgtcgaaag attctgttct tggcttattc aggcaacacg cagataccca tcccgaacgc 60
 ccgcctctcg tcgatogcga gcgctcgctt agctaccgcg aactcgaccg gctcagcgac 120
 cggctggccg cccacctggc caggcgcggc gtcgcccggg gcgagctgct gcccctgctg 180
 gccgaacgct cggccgaact ggtcatcgcc atcctggcgg ccgccaaagt cgcagcggcc 240
 tacgtaccgg tggacgctcg gcaaccgcag aggcgcaagc gggaagtcct ccgccagtgc 300
 caagccccct tggccctcgc caccatgccc gaggacctgc cggggcaacc ggtggaggtc 360
 atcgcacagg cgctcgcgac gagtgcggcg ggtgcgcgcg cgagaccggc gctcgacggc 420
 agcgaagcgc tgtatgtgat ctacacctcg ggcaccaccg gcgaacccaa gggcgtggtg 480
 atcgagagtc gtcctctggc caacctcgtg ggtgggcaca accggcgctt caatatggtg 540
 caacggagcc gcaccacct gatggcggcg gtcggcttcg acgtttccca atgggaaatc 600
 tgggtccacc tgtgcgcagg cgctgcctc cacttggtgc ccgacgaggt gcgcccagac 660
 ccggcgggcg tgcgtggcatt ctgcgcgag cagcggatca gccacgcctt cgcgcctacc 720
 gtgatggtgc ccgcgctggc ggagcagccc gcccgcgct cgctggcgct gcgctacctg 780
 ttctgcgcgg gggaaaaact gccgcgggtc gcaaccggcg ggctgcccta taccgtggtg 840
 gattactacg gcccgacoga ggccacggtc ttgcgccact gccgcacgt cgacgccgaa 900
 gcacatcggc gaccgcctc gatcggcagc cccatcgacg gctgcgaggc attcatcctc 960
 gacgcgcagc accggccttg ccatggcgac cgaccgggtg aactgaacct ggcgggcgtc 1020
 tgctggcggc gcgaatacct gcgcgacccg gacatgaccg ccaggcgctt ccactactcg 1080
 caggcaactg ggcgtcggtc ctaccgcacc ggcgacaagg cccgctggtt ggccgatggc 1140
 agcctgcagt tcctcggtcg gctggacgac caggtgaaga tccgcggcca ccgcgtcgaa 1200
 ctgcgcgacg tcgaggccgc gctgttgccg cagccggcta tccacggcgc ggtggtgctg 1260
 gcgcatgccg acccagctc cggtagccag caattgagcg cttcgtggt ccccgccag 1320
 caggacggcg atgccagggc cgtgctcgcc gccatcaaga ccgcaactgc ccaggaactg 1380
 cccgactaca tgcgcgcag ccgctacctg tcgctggaca gcctgccgac cacggtcaac 1440
 ggcaagatcg accgcaggc cctgcgtcga cactggagc aacaatgcca ggaacgactc 1500
 gacgagcaac gcttcggcac cccggcgcaa ctgcaagtgg cctgtcctg gcaggaagtg 1560
 ctggggcata ccgacttcgg cctggacgac agcttcttcg aggtcgcgcg ccattccctg 1620
 ctggccgcgg cctggtgag cgaattgagc cgacgcttcg gcaaccgtgc ctacatccac 1680

gacatctacc gcaccccgag cgtgcgccaa ctggcggcca gcctggcgcg gcgcgccggc 1740
 gaagcgccgc cggcgctgga cagcgaaccg gccaggagc tgcaacggga cgtgcgcctg 1800
 cccgccgacg tggatttcag ccgccccacg gacaccggcc aattgctggc gccacggcac 1860
 atcctgctca ccggcgccag cgggetgatg ggcgccacc tgctcgccga gctgctggcc 1920
 agccgcgagc ccgacctgca ttgtccggtc cgtgcgcaaa acgacgcca tgccctcgaa 1980
 cgcctgcgcc aggcgcgccg gcagcaccgc atcgaactcg ccgagacgga ctggcgacgg 2040
 gtcagggcct acgcgcgcga cctgcgagaa ccaggtttcg gactaccggc ggaaacctat 2100
 cgcgagctgg ccggcagcgt cgaccaggtc ttccattccg ccagcgcggt gaacttcata 2160
 cagccataca gctacatgaa gcgcgacaa gtcgaggggc tcggccaggc cctgcgcttc 2220
 tgcgccagcg gccgctgcaa gccgctgatg ctgctgtcga gcatctcggc gtacagctgg 2280
 ggccacctgc ataccggcaa gcgcctgatg cgcgaggac acgacatcga ccagaacctg 2340
 ccggcggttg tcaccgacat gggctacgtg cgcgcaaat ggggtgatga aaagatcgcc 2400
 gacctcgccg ccgaacgcgg cctgccgctg atgaccttc gcctcgcta cgccacctgc 2460
 cacagccgta ccggcgccca cgcgcactac cagtgggtga gccggctggc gcggacctgc 2520
 ctggagtacc gggccgtgcc gctcctgcgc gagctgcgcg agggcctgac cacggtggac 2580
 tacatggtag aggcgatcag cgtcactgcc cggcagcctt cggcgctggg caagaaattc 2640
 aacctggtgc cgagcattcc gcgtgcctg accgtggac agttcttcgg ccgtctcgga 2700
 cgacgcgccg ggcgtccctt tcggcagatg ccgttcgacg actgggtaag tctctgggaa 2760
 gacaatcgcg acgccccgct ctatcccttg ctgagcatgt tcgcgacaa catgtacgcc 2820
 ggccgcagca ccgtcgagtt gtaccaggac acctatctct gggactgcac caacgtcgag 2880
 gaacacctgc gcgggagcgc cgtgcgcgag ccggagttcg acgaccgcct gctcgacctg 2940
 tacctcgccg gcctgggcgg cagcgccatg cggtaa 2976

<210> 17

<211> 1092

<212> DNA

<213> Pseudomonas aeruginosa

<400> 17

gtgggacggc ttcgacgagc cgcgctgcac cctgctggag gcgaaggcca actacgcctt 60
 cctgttcgtc ccgctgctcg gcgtgcccag gccctgggca cgggccaagg tgaagtcgga 120
 cctgctgcag aaggccgagg tccacagcga caaggcccga ccgacccgc cgggtgttcg 180
 cgaatggcac ttcttgacg ggatcgtcta cgagtactgc gccgcggagt acctgcgcat 240
 gggactggcc aacctgaagg cattctggaa tccgatgccg ggaacggacg agcacgacga 300
 ctaccaggaa acccgcgcca aggaacagga agagatgaaa aggttttgcg aagagaacct 360
 ggggtattgc gcatgacgga cgccaaggct ttcaggcgct acatattcga gctgtacttc 420
 gatccggcac ggctcctcga actggacgac gaccagcacc tgcaacggat agaacgcttc 480
 ctcgatgccc tcgcgcccct ccatooggcg ctggagaact ggtatctgtg cggcgactcc 540
 ctgcgcgatg cctcagcca caacgtcacc gagcacgcgc aggatctcgc caaggccctg 600
 tcgcgtgacc gacgcacccg ggcgggtgga ctggtgctat ggaacggcga ggaggatccg 660
 ctcaaggcg gggtgtcgct ggactacgag gccagcggca gggccgtctc gtccaggctc 720
 cagttggaag atgcggcag cctgctgcag gtgttcgacg caccggcgct ctccttcgct 780
 gcgatcttcc tcgcggtgct ggaaatctgg cccgaaacga cctggggcat gctcgctccg 840
 catgcgtact tcgtacacca gcggaccttc ccggaccgcc gcagcatcgg ctggatcggc 900
 ttctgcccgc atccgctaag ggccacggac ttcccggcgg ctacggagct ggtcgacatt 960
 cccggccgtg gcacctgct gctgaacggc cgcgaaccga tggacgaaac ccgtcgcgaa 1020
 catttcgagc gcgtcggcga agcggacatc aagctgatgg aactgggcta cctgccgcgc 1080
 ctgcgcggct ga 1092

<210> 18

<211> 1281

<212> DNA

<213> Pseudomonas aeruginosa

<400> 18

atgcacgcca tcctcatcgc catcggtctg gccggcgacg tatttcctt catcggcctg 60
 gcccgacccc tgaaactgcg cgggcacgcg gtgagcctct gcaccatccc ggtgtttcgc 120
 gacgcgggtg agcagcacgg catcgcgctt gtcccgcgtg gcgacgaact gacctaccgc 180
 cggaccatgg gcgatccgcg cctgtgggac cccaagacgt ccttcggcgt gctctggcaa 240
 gccatcgccg ggatgatcga gccgggtctac gagtacgtct cggcgacgag ccatgacgac 300

```

atcgctggtgg tcggctcgct atgggcgctg ggcgcacgca tcgctcacga gaagtacggg 360
attccctacc tgtccgcgca ggtctcgcca tcgacctgt tgtcggcgca cctgccgccg 420
gtacacccca agttcaacgt gcccgagcag atgccgctgg cgatgcgcaa gctgctctgg 480
cgctgcatcg agcgcttcaa gctggatcgc acctgcgcgc cggagatcaa cgcggtgcgc 540
cgcaaggtcg gcctggaaac gccggtgaag cgcattctca cccaatggat gcattcgccg 600
cagggcgtgg tctgctgtt cccggcctgg ttccgcgcgc cccagcagga ttggccgcaa 660
cccctgcaca tgaccggctt cccgctgttc gacggcagta tcccggggac cccgctcgac 720
gacgaactgc aacgctttct cgatcagggc agccggccgc tgggtgttcac ccagggtcgc 780
accgaacacc tgcagggcga cttctacgcc atggccctgc gcgcgctgga acgcctcggc 840
gcgcgtggga tcttctctac cgggcgcggc caggaaccgc tgcgcggtt gccgaaccac 900
gtgctgcagc gcgcctacgc gccactggga gccttgctgc catcgtgcgc cgggctggtc 960
catccgggcg gtatcggcgc catgagccta gccttggcgg cgggggtgcc gcaggtgctg 1020
ctgccctgtg cccacgacca gttcgacaat gccgaacggc tgggtccggc cggctgcggg 1080
atgcgcctgg gcgtgcgctt gcgcgagcag gaggttgcgc gggcgctgtg gcgcttgctc 1140
gaggaccccg ccatggcggc ggctgtcgg cgtttcatgg aattgtcaca accgcacagt 1200
atcgcttgcg gtaaagcggc ccagggtggtc gaacgttgctc atagggaggg ggatgctcga 1260
tggtgaagg ctgcgtcctg a 1281

```

<210> 19

<211> 651

<212> DNA

<213> Pseudomonas aeruginosa

<400> 19

```

atgccgcctt ttttttctcg gccggcacga cagggggact tggctcatgat cgaattgctc 60
totgaatcgc tgggaagggt ttccgcgcgc atgatcgccg agctgggacg ctaccggcat 120
cagggtcttca tcgagaagct gggctgggac gtggtctcca cctccagggt ccgcgaccag 180
gaattcgacc agttcgacca tccgcaaac cgtacatcg tcgcatgag ccgccagggc 240
atctgcggtt gcgcgcgctt gctgcgcagc accgacgcct acctgctcaa ggacgtcttc 300
gcctacctgt gcagcgaaac cccgcgcgag gatccgtcgg tctgggagct ttccgcgtac 360
gccgccagcg cggcggacga tccgcagctg gcgatgaaga tattctggtc cagcctgcaa 420
tgcgcctggt acctgggcgc cagttcggtg gtggcggtga ccaccacggc catggagcgc 480
tatttcgttc gcaacggcgt gatcctccag cgctcggcc cgcgcagaa ggtcaagggc 540
gagacgctgg tcgcgatcag cttcccggc taccaggagc gcggcctgga gatgctgctg 600
cgctaccacc cggaatggct gcagggcgta ccgctgtcga tggcggtgtg a 651

```

<210> 20

<211> 1167

<212> DNA

<213> Pseudomonas aeruginosa

<400> 20

```

atgcctttga ttgtctatgt gctcggtgcc gcgatcttcg ccctgaccac cagcgaatac 60
atggtcgccg ggctgatgcc ggcgctggcc gccgaattcg gctgtcctt cgcgcgcatc 120
ggctacctgg tcaccttcta cgcgcgtgcg atggcgcgtc gggcccgct gttgaccacc 180
gccctgctcc ggggtgcgcg caagaacgcc ctgctcggcc tgatcgcgct gttcgtggte 240
ggccaggtea tcggcgccct ggcgcggg gcgatcggga tggtcgcggc gcgactggte 300
accgcggtcg ccgcgcgcgc cttcttcggc gtggcgctga ccgctgcgc cgaactggte 360
gaaggcaacc agttcggcgc cgcgtcgtcg ctggtgctcg gtggcctgat ggtcggcacc 420
gtgctcggcc tgcccgtcgc cacctggctg gggaatggt acggctggcg cgcgagcttc 480
ttcgcggtgg cgctgggtgc ggtgctggtc ggctgctgg tgttgacgt gatgccggcg 540
atcccggggt cggcgggcag cggctcgctg cgcgaggaa tgaagggtt caggaacgcc 600
catctatggt ggggtctacgc caccagcctg ctgctgatcg gcgccacct cgcggcttc 660
acctatttcg tgccgatcct caccgaggte agcggttct cgcctcgac cgtaccgctg 720
ctgctgggtg tctacggcct ggcgacgctg gtgggcaaca acatcgctcg ccgcctggcc 780
gaccgccata ccacgcgggt cctggccttc ggctgctgg cggccatcgc cgcgatggte 840
gccttcgccc tgttcggaca ggttcggcg gtggcggtgg cggcgctggt ggtgatcggc 900
ctgaccgggg tgtcgatgaa cccggcgctg gtgaccgcgc gcgcacgggt cggccataac 960
aacatgctgg tcaactcggt gcacactgcc tgcatcatgc tcggcgtaac ggccggttcc 1020
tggatcgcgc gcctgggcat cgcggcgga ttggcctgc agggcgcgct ctgggtcggc 1080

```


gcggccctcg gagtactggc gctgctgacc ctgctgccgg agctgcgctt cgcccgcgcc 1140
ccggtaggcg gggcgctggg ccgctga 1167

<210> 21

<211> 993

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 21

atgccgcgcg ccgcgctggt ctggggcctg ggcagctacc tgcccagggc cgtgctcagc 60
aacgacatgc tcgcgcgcga gctggacact tccgacgcct ggatcagcag ccgcaccggc 120
gtgcgccagc ggcataatgc cggcgacctc ggcagcggcg acctggccct gcggggcgcc 180
tccgcgcgcg tcgcctcggc ggggctggag cgagtcgatg cgggtggtgct ggcaaccagc 240
accggcgact tctgctgccc ggccaccgcg cccagggtcg cggcgcgccct ggggttggtc 300
ggcgcgctcg cgttcgacct gtcgcgcgcc tgcaccggct tcgtctacgg cctggccagc 360
gtcggctcgc tgatcagcgc cgggctggcg gacagcgcgc tgctggtcgg ggtggacact 420
ttcagccata cctcgaccc cgcgatcgc tcgacccgcg cactgttcgg cgacggcgcc 480
ggagcggttg tgctgcgtgc cggcgatgcc gaggaagaag gcgcgctgct ggccttcgac 540
ctcggcagcg acggccacca gttcgacctg ctgatgacct ccgccgtcag tcgcgccgaa 600
cgcagttccg gacaggcctc caactacttc cggatggacg gcaaggcagt gttcggccag 660
gcggtgacgc agatgagcga ctcgggtgcg cgggtgctcg accgggtcgg ctggcaagct 720
tcggacctcc atcacctggt ccgcaccag gccaacacac gcattctcgc ggcggtcgcc 780
gaccagctcg accttcccgt cgagcgagtg gtgagcaaca tcgccagggt gggcaatacc 840
gtcgcgcgct cgattccccc ggccctggcc cagggcctgc gccaaaggcat cctgcgcgac 900
ggcggaaca tggctcctcac cggtttcggg gccggactga cctgggggtc ggtcgccctg 960
cgctggccga agatcgttcc gacaatggac tga 993

<210> 22

<211> 1257

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 22

gtgcctgac gcaaactgag actgggcgag gaactgatct cgccactgca cgcgctctac 60
gacggcctgc aggtggacgg cgcgcgcgt cccgcgcac cgcgcgcga gcatccggtg 120
tgggtggtga cgcgctaccg cgacgcgcgc aaggctctca accatccggg cgtccgcgcg 180
gacgcccggc aggcgcgcga actctacgcg aagcgtaccg gcagcccgcg cgcggggatc 240
ggcgagggac tcagccacca catgctcaac ctcgacccgc cggaccatac ccgcctgcgc 300
tcgctggttg gccgcgcgtt cccccgcgc cagggtggag gcctgcaacc gcatatagaa 360
cggatcaccg aggcattgct ggacgccatg gccggcgcg aacaggccga cctgatggcc 420
gacttcgcga tcccgtcgac catcgcggtg atcttcgagc tgctgggcat tcccagggcc 480
gagcgcgaac acgcccgcga gtccctggag cgcagggcg aactgctgtc gccggaggag 540
gcccaggccc tggccgatgc gcaggtcgac tacctgcgcg tgctgctcga ggccaagcgc 600
cggcagccgg ccgacgacgt ctacagcggg ctggtgcagg ccgcgcgca gagcggccag 660
ttgagcgaag cggaactcgt ctccatggcc cacctgctga tgatgagcgg cttcgagacc 720
accatgaaca tgatcgccaa cgcgctggtc acctgctgg tcaaccgcga gcaactggcg 780
ttgctgcggg cgcagccgga actcctgccc aacgcctagg aagaactggt ccgccacgac 840
agtccggtgc gcgcctcgat gttgcgcttc accgtggaag acgtggaact ggacggggtc 900
accattccc cgggcgaata catcctggtc tccaaactga ccgccacca cgatgccgag 960
cgcttcgacg atcccgaccg cctcgacctc acccgcaaca ccgatggcca tctcggtac 1020
ggcttcggcg tgactactcg cgtcggcgcc tcgctggccc ggctggaggg gcggatcgcc 1080
atccagcggc tgctcgcgcg ctcccccgc ctccagttgg cggtgcccca cgcggagctg 1140
cagtggctgc cgatcacctt cctccgcgcc ctgatcagcg tgccggtgcg caccggatgc 1200
agcgcgcccg cgaacaccgc ctcccacgcc aaccgcgatc agaggatcgc ccaatga 1257

<210> 23

<211> 915

<212> DNA

<213> *Pseudomonas aeruginosa*

2025 RELEASE UNDER E.O. 14176

<400> 23

```

atgttattca ccagcaaacc tctctcgccc cagggccgccc acgtactgat caccggcgccc 60
tocagcgccc tcggccggga aaccgcgctg cacctggccc aacagggttt ccaggatgatc 120
gccgggggtgc gccgccagga ggatggcgag cgcctggcga acgcctgccc gtccggcgccg 180
atcagcacgc tgctgatoga tgtcaccgac gaggaatcca ttggccgggc cgccgcgcag 240
gtggcggaga aagtcggcga taccgggctc tggggcctgg tgaacaacgc cgggatctgc 300
atttcgcgcg cgctggaatg cgtctccagc gacctgctgc ggcgccagct ggaagtcaac 360
ctgatcgccc agctcgcggt gaccggggcg atcctgccgc tgctgcgcgc tggcggcgcg 420
gcgcgcctgg tgaacgtcac ctcgggcctc ggctcggtcg ccattcccta cctggggccc 480
tactccgccc cgcagttcgc caaggaggga gtgagcgacg ccctgcgcgc cgagctggca 540
cccatgggca tccaggtctc ggtggtcagc cccggggcga tctggacgcc gatctggggc 600
aagatcgcca gcgagggcga gcgcgcctg gccgacgccc ccgacgccgt cgccgacctc 660
tatcgcgata cctacctgcg ctctctccag gccaacgagg acggcgcgcg caacagcgcg 720
accaagcccc cgatgtcgc cgccgcggtg catgccgcgc tcaccgcggc caagcccgcg 780
acccgctacc gggtcggcgc cgacgtgcgc cgcggtaccc tgctggcgcg gctgctgccc 840
gatagcgtga tcgacgggat gttccgcccc atcgtcaccg ccgccccggc ggcgaaggag 900
gagcaacgtg cctga

```

<210> 24

<211> 1329

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 24

```

atgatggccc agatacgacg cccgctgtcc gcgggtggaac gctgggtactg gctcagcgac 60
cagttctccc cgctgaacgt gatttcccgg gtgcgggtcc atggccggtt gtccatcgac 120
gacctgcgcc cgggcctcga cgcgctgcag gcgcggcatc cgctgctgcg cgcgcggtac 180
gagcacgatg ccgggctcga tccgcgctgg gtgccctgcg agcggcccat cccgctgcgc 240
gaggtgcgcg gcggcggcga ggagcaatgg ctgcgggaaa tcaacgagcg cgaattgccg 300
gaacgcacat atccggacag cgggccactg atccgtaccg tggcgatcgc caccgacgcc 360
ggcgccccac acctgctggt cgtgggtaccg cacatcatcg ccgacggcac taccgtgctg 420
accctcgccc aacaatgggt gaccctggcc gccgaccccg ccgcgcaacc ctggaccgcc 480
agcgccctgc cgccggcgga ggatctgcgt ccgcgcgcgt tcaccggcga cgaaggcgcg 540
gcgcgcctgg ccgagcagac cgcccaggac gaagcgctgg tcggccgcca ccgcccgggc 600
cggatcgagc cgagcaaccc ggtgccgctg gaagcgcggc gtaccgcct gctgcaccgg 660
gagctggacg gcgcgcagct ggaacagctg caacgacgcg ccgcgaaca cggcaccacg 720
gtacacggcg cgctgaccgc ggcgctggcc atcgccgcgc gccacgacca ccagcgccgc 780
cctagccaca tcgccatogg ctgcgcgac gacttccgcg acgaactgga gccgcccgtg 840
cgccccgacg aagtaggcac ctacgtcgcc acggtaccgg tgggtgctgga catcgcccgg 900
ccgttctggg aggtcgcccc cgcgctcacc gacgacctcg gcgaacgccg tcgccagggc 960
catcatttca acctggtcac cctggctgcc agcgctgcgc cgcgctgcat ggccgacgcg 1020
cggccattca tggccttcac ggaagccgaa gggccgatca acctgtgctc ctccaacatc 1080
ggtoctatc cgttcccga cgggatcggc gccttgccgc tctccgacgc gcagttcctc 1140
accggcatct cggtagaacg ctacttcgtg gccgccatca actccagcca tggccggctg 1200
ttctggaact tcacctatat cgacgaagcg gtccccggcg aacgcgccga acgcctggcc 1260
gaagattgcc tgggcaccct gctgtcggcg atccacgccc cccaacgatc cgccctcgag 1320
gagcaatga

```

<210> 25

<211> 1167

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 25

```

atgagcagac atccccctgaa gatcgctcacc gccggcgccc gcacgcggcg gctcgccgcg 60
gccgcctgcc tgaagaccgc cggcttcgag gtccgaactct acgagcgggc caggagagctg 120
cgcgcggtcg gctcggcgct gtgcgtgatg cccaacgcgc tgaccgccct ggagagggtc 180
ggcgctgcgc ctgaccttac ccgcgcccag gccttcgact cgctgcgggt cctcaccgcg 240
cgcgggcgac cgatccgcgc catcgacttc ggcggcctgg ccgctcagct cggccagccg 300
agcctggcga tccaccgcgc gagcctgcag caggcgctgc tgggaacagg ccgcgactgc 360

```



```

cgcgacgacg cgcgcctgcg cgcactgccc ctggagcacg ggacgaacat cgagtacctg 6000
gccccgcgcg acgactacga ggcgaccctg gccggactcc tcggcgagtt gctggatcgt 6060
ccccgggtag gcaccccgga cagcttcttc gacctcgcg gcacctcgct cagcgcgatg 6120
cgcttcatgc tgctgatcga gaagcgctat ggcgtcgacc tgccgatggc cgcgctgac 6180
gagacgccga ccgtggaggg cctggccgaa cgctgcggg aacgctcggc ggtgcgcgcc 6240
ttcgacccgc tggtagcgat ccgtgcccgc ggcagccgac cgccgctgtt cctcgctcc 6300
ccgctcgggc gccacgtgct ctgctacctg ccgctgggtc gcgactgcc gccggaccag 6360
ccggtatatg ccctgcaggc ggccggcacc ggccagggca gtacgccgct ggcggtcctc 6420
gaggacatcg ccgccagtta cctcgcgggc atccgcccgg tgccagccga agggccctat 6480
tacctcgggc gctggtcggt cggcggtctt gtgcctacg agatggccc gcaactgcgc 6540
gcgctcgacc cgcaggcggt cgcaccaact atcgtgctcg actccatcac cgtcgaccgc 6600
aaccacgccg gcagcgccag cgacgaagcc ctgctgctgt tcttctactg ggaactggtc 6660
tggttcgagc gcagcgacaa ggaggtcgag ccgctgcctg aaggcgcgag cctggagcag 6720
aaactcgacc acatcgctga acgcgccatc gaggccggcg tacttcccgc cggcaccgcc 6780
cgcgccaccg tgcagcggtc ctacgagctg ttccggggcg gctggcaggc actcatcggc 6840
tatcgccggc aagtcagcga ccaggacatg accctgctgc gcgcgacgg cccgctgcgc 6900
ctggcgctga agccgatgca cgacgcccgc ggcacccact acggcgaccc gaagaacggc 6960
tggcagcact ggaocagtgg ccgcctcgat gtgatcgacg tccccggcga ccacctgggt 7020
ctgatgaaag aacctatgt cgagacggtc gcggcagaga tcgcccgtt gtcgaaccc 7080
tccacctcca gcgaacggac ccgcccatga 7110

```

```

<210> 27
<211> 1404
<212> DNA
<213> Pseudomonas aeruginosa

```

```

<400> 27
atgaaaacgc ccgctggac ggcacatgcc ctctgggtca tgccgctcgc cctggggctg 60
caatccgcgc tggtagcggg ggatgagcag ccaagcaaga cttccagcta ttccgcggtg 120
gtgatcaatg aggacttcgc caccatcatg aagcgcatga cggcgaacaa accgtcgatc 180
gaacaggccc acaagacgct tctcgagcag cgttacgac tccagcgacag gccggccaag 240
ggcgccagca tgacgcgcgg caagccgctg caggagggga tccgggtgaa gctgccggcc 300
ggcaccagct gggaggaaact ggccaggctg agccccgagg aaatccgcaa gcaggggctg 360
ttcccggtg gcttccctgc gctgcgcac cccaaccatg ccgaaggcgg gatggtcttt 420
cccaagttcc tcacgacga gatcaagcgc caggaaaagc gcgacctgac ccgtttcgac 480
ctcgactacg acctgcgcga ccacttctct cgggaattcc cggcaccgat gttccttacc 540
accggcgctg acctgggcga tgtgtccaag ggcaagctgg tgaccatcga caactatttc 600
gagttgttca acgggattct caatcccaag cagctggaag ggctgcgcct gctgctaacc 660
gcctttccgc agcagcagtt caacctcacc gacgatcgcc gtagcgagca tccgagccgc 720
ggcgtagcct gcttcgactg ccacgcgaac ggccacacca atgccgctac tcacctggcc 780
ggcgatgtgc gcccgcagcc gttccgccac cgcacgcaga caccgacgct gcgcggggtg 840
aacatccagc ggttgttcgg ctgcagagg gcgctgagga cttaccaggc cttaccaggc 900
ttcgagcagc gcgcgccta cttcgacggt gatccggtaa tcgccaccaa gaagggggtg 960
aacgtgctcg agcgtggcag tcaagtgcac ttcattgggt agttccaggc gctgctggac 1020
ttcccccccg caccgaagct ggatgtggag gggcggtctg atccgggcaa ggccagcgag 1080
caggaaattg gtggcgaaaa gctgttctac ggcaaggcgg cctgcgcggg gtgccatgcg 1140
ccgccttact tcaccgacaa cctgatgcac aacctgaagg tggagcgctt ctacgatccg 1200
aaactgggtc atggcggtgat ggcgccgcc gacgggcca tcaagacctt cccgttgcgc 1260
gggatcaagg attcgccgcc gtacctgcac gacgaccgcc tgcagacctt ggaggacacc 1320
gtggagttct tcaacctggt actggagcgc aagctgtccg cggaagagaa gggcgacctg 1380
gtggcctacc tgcgtacctt gtga 1404

```

```

<210> 28
<211> 1386
<212> DNA
<213> Pseudomonas aeruginosa

```

```

<400> 28
atgctcacgg tgtgtgcgaa ccccaagggt atttcccgac cagctcccc gcggtcggga 60
tttttttttg cctgtcgctc agcgcttcgg gtccgaaggc gaatagcccc gccggcgag 120

```

2025 RELEASE UNDER E.O. 14176

gctcgccagg ggcgcgaaca ggggctccgg cagatcggcc caatcgaacc agggccagcc 180
 gtcgcacttg tccggctcca tgaggcgcg ctcggcatcc tccgcgcaac cggccaggat 240
 gaacgcggtg aggtagtggc gcccctcgaa gacgtcattg ctgaacgggc cgtggcgag 300
 ttcgctcagc gccaggctcg tctcttccag ggcttcgcgc agggcgagc cctccaccgc 360
 ctcgcggaac tcgagatggc cgccgggcg cgaccagcag ccagcgccat gactgccctt 420
 gcggcgcccc agcaacacct tgccgtccc caagatcagg acgcccacgc ctacctgcgg 480
 tgccggcatc gtcgtactcc tgcttcggga tcagagatgg agcgtaccgc tcatgtacaa 540
 cgccgccttg ccggagatga ccaccgctc gccgcgcacg tcgcattcca ggcccccctt 600
 gcgcgccccg ccctgctcgg cgctcagccg ggtcttgccc aggcgctgcg ccagtacagg 660
 cgccaggagg gtatgcgcgg agccggtcac cgggtcttcg ttgacgccga cgttggggcc 720
 gaaccagcgc gagacgaaat cgaagcgctg gctgcgcgcg gtcaccgcca ccccgcgga 780
 cggcaagccc ttcagccggg cgaagtcagg cgccaggggc gcgatcgtct tttcgtcgtc 840
 gaccaacca aggtaatcgt cggctcttcag cacttcgcgc tcggcaatac ccagcgctc 900
 cagcagtcgg tccggtgtcg cgcaaggctc cggacgcttg gccgggaagt ccacgccag 960
 cgagtcgccc tcgcgcgcga cgctcagctc accgctacgg gtacgcgaaac gcagtaccgg 1020
 ggaagcgtcg tcgagcttgt ggatcagtag ccaggcgctc gccagggtcg catgaccgca 1080
 caggtccacc tcgacctgcg gcgtgaacca gcgcaatcga tagtcgccg cgcgccgac 1140
 gacaaaggcg gtttccgaaa gattgttctc ttccgcgatg gcctgcaggc gctcgtcgtc 1200
 cagccaggca tcgagggggc agaccgcgc cgattgccc tggaaggggac tgcggcgaa 1260
 tgctgtacc tggaagatcg tcagttccat gttccggact cctgtatcga tgggctgcgc 1320
 accttagcag ccggaccgag accaggacaa tgccgcgccc cgcgaggcg cctcgctcag 1380
 atctga 1386

<210> 29

<211> 1104

<212> DNA

<213> Pseudomonas aeruginosa

<400> 29

atgaaaaaag tttgtgcaact ggcgttatcg atcctgacga cgatcggtgc gacagcggcg 60
 gacagtgcac gggctgcgca aaccagcgct catctttaca actggtatga cttcatcgcc 120
 ccggaaacgc ccaaggcttt ccagaaggaa accggcaccc gtgtcgtcct cgacaccttc 180
 gacagcgccg agaccgcgca gggcaagctg atggctcgcc gctccggcta cgacgtgggtg 240
 gtgatcacct ccaacatcct gccgggctg atcaaggcgg gcgtcctcca ggaactcgac 300
 cgcgaccggc tccccactg gaagaacctc gacgcggaca tcctcgggaa gcttcaggcc 360
 aacgatcccg gcaatcgota tgccgtacct tatctctggg gaaccaccgg gatcgctac 420
 gatgtggaca aggtccgcaa gctgctcggc cccgacgcgc cggctcgact ctgggacctg 480
 gtcttcaagg aggagaacat ctcccgctc agccagtgcg gcgtggccac gctggactcc 540
 tccaccgagc tgggtgtccat cgccctcaac tacctgggccc tgccgcacaa cagccagaat 600
 cccgaggact accagaaagc ccaggaactg ttgctgaagg ttgcccccta cattcgctat 660
 ttcgactcct ccagagtcga caccgatctc tccaacggca acgtctgcgt ggtggtcggc 720
 tggcagggca cggcctacat ggcccagggt aacaacgaac aggcgggaa cggtcgccat 780
 atgcctaca gcattcccc ggaaggctcg ctggtctggg ccgagaacat ggtgctgctc 840
 aaggatgcac cgcatocgca gcagggttat gcgctgatcg actacctgct gcgtccggag 900
 gtcacgcgca ggacctcaa ctacgtgggc tatccgaatg gcaaccaggc ggcgctgccg 960
 ctggtagagc ggaaactgcg ggaaaacccg gcggtttacc tgagcaagga aaccatggcg 1020
 accctcttcc cgctggaaac cctgccactg aaggctcgaga gaatccgtac ccgggtctgg 1080
 agccgggtca agaccgggag ctga 1104

<210> 30

<211> 1251

<212> DNA

<213> Pseudomonas aeruginosa

<400> 30

gtgggctgtc cggggcggtt aggatggaca ttttcatcgt ctcgggcagg cctgtcgcga 60
 ccgcgcgaag tcgcgacgga tgccgctgct aaggagcaac ggatgaccgt tcttatccag 120
 ggggcccggg tcgcgggccc ggcgctggcg cgcgaattca ccaaggcagg catcgactgg 180
 ctgctggctg agcgggccag cgagatcagg cccatcggtc ccggcatcac cctggcgagc 240
 aatgcgttga cggcgttgtc cagcaccctg gatctcgacc ggctgttccg ccgtggcatg 300

```

ccgttggcgc  gcatcaacgt  atacgcccac  gacgggttcga  tgctgatgtc  gatgccttcc  360
agtctgggtg  ggaattcccc  cggcggcctg  gcgttgccagc  gccacgaact  gcatgcggcg  420
ctactggagg  ggctggatga  gtccgcgcat  cgggtcgggg  tctccatcgt  gcagatcctc  480
gacggactcg  accacgaacg  cgtgaccctg  agcgacggca  ctgtccacga  ctgttcgctg  540
gtggtcgggt  cggatggcat  tcgttcgagc  gtgcgacgtt  atgtctggcc  ggaggcgacc  600
ttgcgtcatt  ccggcgaaac  ctgctggcgc  ctggctcgtt  cccatcggt  ggaggacgcc  660
gagctggcgg  gagaggctct  ggggcacggc  aagcgctcgt  gcttcatcca  gatcagcccg  720
cgcgagatgt  atgtctacgc  gacctgaag  gtgcgcgggg  aggagcccga  ggacgaggag  780
ggcttcgtaa  ccccgcaacg  gctggccgcc  cactacggcg  acttcgacgg  catcggcgcg  840
agcatcgccc  ggctcatacc  gagcgccacc  acgctgggtc  acaacgacct  cgaggagtgt  900
gcgcggcct  cctggtgcgc  cggacgggta  gtgctgatcg  gtgacgcgc  acacgccatg  960
acgcgaacc  tggggcaggg  cgcggccatg  gccctggagg  acgccttct  gctggcgcg  1020
ctgtggtgc  tggcgccgc  cgccgagacg  ctgatcctgt  tccagcagca  acgcgaggcg  1080
cggatcgagt  tcatcaggaa  gcaatcctgg  atcgctcgcc  gccttgggtc  gtgggaatcg  1140
ccctggagcg  tctggtctag  gaataccctc  gttcgccctg  tgccgaatgc  cagtcgcagg  1200
cgctccacc  agcgtctttt  caccggtgtc  ggtgagatgg  ccgcacagta  g  1251

```

<210> 31

<211> 1754

<212> DNA

<213> Pseudomonas aeruginosa

<400> 31

```

atgatggacg  ccttcgaact  tcccaccacc  ctgggtccagg  cctgcgctcg  ccgcgctgtc  60
caggagcccg  agcgccctgg  gctgcgcttc  ctgcgcgagg  acgatggcga  aggcgtggtc  120
ctcagctatc  gcgatctcga  cctgcgcgcg  cggagcatcg  ccgcggccct  gcaggcccat  180
gcgcagctgg  gcgatcgcg  ggtactgctg  tttccagcg  gcccgcacta  cgtcgcgcg  240
ttcttcgggt  gcctgtatgc  cggggtcctc  gcgggtgcgg  cctaccgcgc  ggaatcggcg  300
cgccgccatc  accaggaacg  cctgttgctg  atcatcgccg  acgcgagcc  gcgcctggtc  360
ctgaccaccg  ctgacctgcg  cgagccattg  ctgcagatga  acgcgcaact  gtccgcccgc  420
aacgcccgc  aactgctctg  cgtcgaccag  ttggaccgg  ccgttgccga  ggcctgggac  480
gagccgcaag  tgctgcgcga  gcacatcgcc  ttctccagt  acacctccg  ttcaaccgca  540
ttgcccaagg  gcgtgcagg  cagccatggc  aacctggtcg  ccaacgagg  gctgatccgc  600
cgaggcttcg  gcctcggtgc  cgacgacgtg  atcgctagct  ggctgcccgt  gtaccacgac  660
atgggcctga  tcggcgccct  gctgcaaccg  atcttcagcg  gcgtaccctg  cgtgctgatg  720
tcgcgcgct  acttcctcga  acgtccgggt  cgctggctgg  aagccatcag  ccagtacggc  780
ggcaccgtca  gcggcggtcc  cgatttcgcc  taccggctgt  gcagcgagcg  ggtcgccgag  840
tcggccctgc  agcgtctcga  cctgagcgg  tggcggttag  ccttctccg  ttccgagccg  900
atccgccagg  acagcctgga  acgcttcgcc  gagaaattcg  ccgccagccg  cttcgacgcg  960
tccagtttct  tcgcctgcta  cggcctcgcc  gaggcgacc  tgcttcgtac  cggcgcccg  1020
cgcgccagg  gcattcccgc  cctggcggtg  gatggcgagg  cgctggcgcg  caaccgcac  1080
gccgaaggcg  aaggcagcgt  gctgatgtgc  tcggcccgca  gccagccgga  acacgccgtg  1140
ctgatcgctg  acgcggcgag  cggcgaggtc  ctgcgcgacg  acaacgtcgg  cgagatctgg  1200
gccgcgggg  cgagcatcgc  ccacggctac  tggcgcaacc  cggaagcttc  ggcaaggcc  1260
ttcgtcgagc  gtgacgggg  cacctggctg  cgcaccggcg  acctcggtt  cctccgcgac  1320
ggcgaactgt  tcgtcacccg  gcgcctgaag  gacatgctca  tcgtccgcgg  ccacaacctc  1380
tatccgcagg  acatcgaacg  caccgtcgag  agcgagggtc  cgtcggcgcg  caagggcagg  1440
gtcgcgccct  tcgcggtcac  ggtcgatggc  gaggaaggca  tcggcatcgc  cgccgagatc  1500
ggtcgcgcg  tccagaaatc  ggtgcgggcc  caggagctga  tcgactcgat  ccgccaggcg  1560
gtggccgagg  cctaccagga  agcgccgaag  gtggtggcgc  tgctcaatcc  cggcgccctg  1620
ccgaagacgt  ccagcggcaa  gctgcaacgt  tccgcctgcc  gcctgcgcct  ggaagacggc  1680
agcctggaca  gctatgcgt  gtttcccgcc  ctccaggccg  tgaggaggc  gcagcccgcc  1740
gcaggcgacg  acga  1754

```

<210> 32

<211> 7335

<212> DNA

<213> Pseudomonas aeruginosa

<400> 32

gtgttggtca	tcacccagca	ccatatcgtg	tccgacgggt	ggtcgatgca	ggtgatggtc	60
gacgaactgc	tccaggccta	tgccgcggcg	cgccgcggcg	aacaaccgac	gctggcgcca	120
ttgacgctgc	agtacgccga	ctatgctgcc	tggcatcgcg	cctggctgga	cagcgcgag	180
ggcgcgcggc	agctggatta	ctggcgtag	cgccctggcg	ccgagcagcc	ggtcctggaa	240
ctgcccgcgc	accgggtgcg	cccggcccag	gccagcgga	gcgggcagcg	tctggacatg	300
gcgctgcgcg	tgtcattatc	ggaggagctg	ctggcctgcg	cccggcgga	gggtgtcacc	360
ccgttcatgc	ttctattggc	ctcgttccag	gtgctgttga	agcgctatag	cgggcagtgc	420
gacattcgcg	tcggggtacc	tatcgccaac	cgcaaccgcg	ccgaggtcga	gcgcctgac	480
ggctttctcg	tcaataccca	ggtgctgctg	tgccaggctg	atgctggcct	ggctttccgc	540
gatctactgg	gccgcgtgcg	cgaggcgcg	ctgggcgcgc	aggcgacca	ggatctgccg	600
ttcgagcaat	tggtcgatgc	cttgagccc	gaacgcaatc	tcagccacag	cccgttgttc	660
caggtgatgt	ataaccacca	gagcgcgag	cggcaggatg	cccaagtcca	tggtttgac	720
atcgagagtt	ttgcctggga	tgggtgctgc	gcacagttcg	atcttgccct	cgataacctg	780
gaaaccccgc	acggccttgg	ggcgcgctg	acctacgcga	ccgacctgtt	cgaggcgcg	840
accgtcgagc	gcatggcgcg	gcattggcag	aacctgctgc	gcggcatgct	ggaaaacccg	900
caggccagcg	tcgactcgct	gccgatgctc	gatgccgagg	agcgtggcca	gttgctggaa	960
ggctggaacg	ccactgccgc	cgagtaccgc	gtgcaacgcg	gcgtgcaccg	gttgttcgag	1020
gagcaggtcg	agcgacgcgc	gacggcgcg	gcgctggcct	tcggcgagga	acgcctggac	1080
tacgcccagc	tgaaccgcgc	ggccaaccgc	ctggcgcatg	ccctgatcga	gcgcggggtc	1140
ggtgcggacc	gcctgggtgg	cgtggccatg	gagcgttcca	tcgagatggt	cgtggccctg	1200
atggcgatcc	tcaaggccgc	cggcgcctac	gtgccggtgg	acccggagta	ccccgaggag	1260
cgccaggcct	acatgctgga	ggacagcggc	gtgcagctgc	tgctcagcca	gtcgcacctg	1320
aagctgcgcg	tggcgcaagg	cgtgcagcgc	atcgacctgg	accaggccga	tgctggctg	1380
gaaaaccatg	ccgagaacaa	tccggggatc	gagctgaacg	gcgagaatct	tgcttatgtc	1440
atctacacct	ccggtctccac	cggcaagccc	aagggtgcgc	gcaaccgcca	ttcggcgctg	1500
agcaaccgct	tgtgctggat	gcagcaggcc	tacggcctgg	gcgtcggcga	cacggtgttg	1560
cagaagaccc	cgttcagctt	cgacgtgtcg	gtctgggagt	tcttctggcc	gctgatgagt	1620
ggggcacggt	tgggtgggtg	cgcgccgggt	gaccatcgcg	acccggcgaa	gctggtggcg	1680
ctgatcaacc	gcgaaggggt	cgacacgctg	cacttcgtgc	cgctgatgct	gcaggccttc	1740
ctgcaggacg	aagacgtcgt	ctcctgcacc	agcctgaaac	gcacgttttg	cagcgcgag	1800
gcgctgtcgc	cggaacccca	gcagcaggtg	ttcgccaagc	tgccgcaggc	cgccctctat	1860
aacctctatg	gcccgaacga	ggcgggccatc	gacgtcacc	actggagctg	cgtggaggag	1920
ggcaaggacg	cgggtgccgat	cggccggcg	atcgccaacc	tgggctgcta	catcctcgat	1980
ggcgacctgg	agccggtgcc	ggtggcgctg	ctcgcgagc	tgtacctggc	cggtcggggc	2040
ctggctcgtg	gctaccacca	gcgtccgggg	ctgactgcgc	agcgtttcgt	cgccagcccc	2100
ttcgtggctg	gggagcggat	gtaccgcacc	ggcgacctgg	cgcgctaccg	cgccgatggg	2160
gtgatcgagt	acgcggggcg	gatcgaccac	caggtgaagc	tgcgcgccct	gcgcacagag	2220
ctgggcgaga	tcgaggcgcg	cctgctggag	catccgtggg	tgcgcgaggc	ggcggtgctg	2280
gcggtggaca	gcaggcagtt	ggtcggctac	gtggtgctgg	agagcgaggg	cgccgactgg	2340
cgcgaaagcg	tggccgcgca	cctggcgaca	agcctgcccg	aatacatggt	gccggcgcg	2400
tggctggcgc	tggagcggat	gccgctgagt	ccgaacggca	agctggatcg	caaggcgctg	2460
ccgcagccgc	aagctgctgc	ggggcgagcg	catggttgcg	cgcagaatga	aatggagcga	2520
cgtatcgcg	ccgtctgggc	ggacgtgctg	aagctggagg	aggtgggcgc	caccgacaac	2580
ttctttgccc	tgggtggcga	ttccatcgct	tcgatccagg	tgggtgagtgc	atgccgtgcg	2640
gcgggcatcc	agttcactcc	gaaggacctg	ttccaacaac	agaccgtaca	ggggctggcg	2700
cgagtgcgcc	gcgtaggggc	tgcggtgcaa	atggagcagg	ggcctgtgag	cggcgagacg	2760
gtgttggttg	cgttccagcg	gttggtcttc	gaacagccga	ttcccaatcg	ccagcactgg	2820
aaccagtcac	tgctgttgaa	gccgcgcgag	gccctgaatg	cgaaggcact	cgaagcggcc	2880
ttgcaggccc	tggttgaaca	tcacgacgca	ttgcgtctgc	gcttccatga	aacggacgga	2940
acctggcatg	ccgaacatgc	cgaagcaacg	ctgggcgggtg	cgctgctctg	gcgtgccgag	3000
gcggtggacc	gacaagcgct	ggagtgcctc	tgcgaggagt	cgacgcgag	cctggacctg	3060
gccgacggcc	cactgttgcg	gagcctgttg	tgagatatgg	ccgacggcg	ccagcgtctg	3120
ttgttggtga	tccaccatct	ggtggtggac	ggggtgtcct	ggcgcatctc	gctggaggat	3180
ttgcaaaggg	cttaccagca	gagcctccgt	ggagaagctc	cgcggtgcc	tggcaagacc	3240
agcccgttca	aggcctgggc	cggccgagtg	agcgagcatg	cccgtggtga	gtcgatgaag	3300
gcgcaattgc	agttttggcg	cgagctgctg	gaaggtgcgc	cgcccgagct	tccgtgcgag	3360
catccgcaag	gcgctctgga	gcagcgttcc	gtacacctcg	tcgagagtcg	cttcgaccgc	3420
agcttgaccg	aacgcttgct	gaagcaggcg	ccggcagcct	accggaccca	ggtcaacgat	3480
cttctgctga	ccgccttgcc	gcgagtggtc	tgcggttggg	gcggcgccct	ttcaagcctg	3540
gtacagctgg	aagggcacgg	gcgcgaggag	ctgttcgcgc	atatcgacct	gagtcgcacc	3600

2025 RELEASE UNDER E.O. 14176

gagctgccgc tggcggcgct gttccagacc gagtcgctgc aagcctatgc cgagcttgcc 7260
 gggcgcgaga cttccagcaa tgacaccgat ttcgatgacc ttcgtgaatt catgagcgaa 7320
 ctagaggcga tctga 7335

<210> 33

<211> 2556

<212> DNA

<213> Pseudomonas aeruginosa

<400> 33

atgctttcca atccaaacct ggacctcgtg tcccgcttcg ttcgcctgcc tctggcgag 60
 cagaaattgt tctatcagcg tgtccaggcc aagggcatga gcttcgcccg cctgccgatc 120
 ccgcagactc gccaggagat ggacaacctg ccgctgtcct atgcccaaga gcggcagtgg 180
 ttcctctggc agctggagcc ggagagttcc gcctaccaca ttcctaccgc cctgcgctg 240
 cgcggcagggt tggacattgc gtccttgacg cgcagcttcg cggcgctcgt cgagcgccac 300
 gaaagcctgc gcacgcggat cgcgcggatg ggtgatgaat gggcgagggt cgtctccgcc 360
 gacgtctcgc tggcgctcga agtcgaagt caacggggac tcgacgaaca gcgattgctg 420
 gagcgggctg aggcggagat cgcacgaccc ttcgatctcg aacagggacc gttactgcgg 480
 gtgactttgc tggaggtgga cgcgcagcag catgtgctgg tcatggtcca gcaccatc 540
 gtctccgacg gttggtcgat gcaattgatg gtcgaggaac tgggtccagct ctatgccgcc 600
 tatagccaag ggctcgacgt ggtggtgccc gccctgccga tccagtacgc ggactacgcc 660
 ctgtggcagc gcagctggat ggaggcgggg gaaaaggagc gccagttggc gtactggacc 720
 ggctgctgg gcggcgagca gccgggtgatc gagttgcccc tcgatcacc gcggcagccg 780
 ctgcgagctc atcgtggagc gcaattggac ctggagctgg agccacacct ggcccttgcc 840
 ttgaaacagc tgggtcagcg caagggtgtg accatgttca tgcgtgtgct ggcttccttc 900
 caggcgctgt tgcctcgcta tagcggacag cgggatatcc gtgtcggcgt gcctatcgcc 960
 aaccgtaacc ggggttgaac cgagcggtcg atcggtttct tcgtcaacac ccaggtgctc 1020
 aaggccgaca tcaatggccg gatgggtttc gacgagttgc tggcccaggc ccgccagcgc 1080
 gcgctggagg cacaggctca ccaggacctg ccgttcgagc aactgggtgga ggctttgcag 1140
 ccggaacgca gcctcggcca caaccggtt tccagggtca tgttcaatca ccaggccgac 1200
 tctcgttcgg ccaaccaggg cgtgcaactg ccaggcctgt cgtgagagcg gatggagtgg 1260
 cggagcagct ccgtggcctt cgacctgacg ctggacgtgc acgaggccga ggacggtatc 1320
 tgggcatcgt tcggctatgc cacggatctg ttcgaggcct cgaccgtcga gcgcctggct 1380
 cggcactggc agaattctct gcgcggcatc gtggccgaac cgggcccggc ggtcgccgag 1440
 ttgccgctgt tgcgtggaag ggagcgcat tgcctgtcgc gggcctgggc agagaacgcc 1500
 gacgagggtg ggttgccgcc cctggtccag ttgcagatcc aggagcaggc ccgtctgcgt 1560
 ccgcaggcgc aagcactggc gctggagggg caggccttga gctacgccga gctcaacgcc 1620
 cgcgccaatc gtctggctca ctgcctgata gcgcgtggcg tcggtcccga tgtgctgggt 1680
 ggaatcgccg tcgagcgctc gctggacatg gtggtcggtc tgcgtggcat cctcaaggcc 1740
 ggtggtgcct atgtgccgt ggacctgacc tatccgcagg accgtttgcg tcacatgctc 1800
 gaggacagcg ccgtcggcct gttgctcagc caggagcatt tgcgtcccgg gctgcctttg 1860
 cacgaaggcc tggaggtgct ctccatcgac cgcctggaac gggacgcac ggtgtctacg 1920
 gatgatccgg tgggtgaact gcggccggag aacctggcct atgtgatcta cacctccggc 1980
 tccaccggaa aaccgaagg cgtggccatc agccatgcgg cgcttgcgca gttctcgcgt 2040
 atcgccagtg gttattccgc gctcaccctg gaggatcgga tattgcagtt cgccaccctg 2100
 agcttcgacg gcttcgtcga acagctctat ccggcgctga cccgtggtgc ctgcgtggct 2160
 ctgcgtggcg gcgacctctg ggataccggt gagctgtatc ggcagatagt cgagcagggc 2220
 gtgacctttg ccgacctgcc caocggctac tggaaacctg tctgtctcga tgccctggcc 2280
 gagccacggc gttcctacgg tgccttgogg cagatccaca tcggtggcga agccatgcca 2340
 ctggaggggc cgaagctctg gcggcaagcc ggcctggggc gggtagggtt gctcaatacc 2400
 tatggaccga ccgaggccac ggtggtgtcc agcgtcttcg attgttccgc cgagaacgcc 2460
 cgggtgggca atgccagtc tatcgccag gcgctaccgc gccgtacgtt gctggtgctg 2520
 gatgaacatc tcggcctact gcccgtaggg cggtag 2556

<210> 34

<211> 2334

<212> DNA

<213> Pseudomonas aeruginosa

<400> 34

atgtcccggc cgttccggcc accactttgc agagaaacga catcgatggg gatgcggtacc 60
 gtactgaccg gcctggccgg catgctgttg ggttcgatga tgccgggtcca ggccgatatg 120
 ccgcgccga cggggtggc cgcggatata cgtggaccg cctatggcgt gccgcacatc 180
 cgggccaaag atgagcgcg cctgggctat ggcacggct acgcctacgc gcgcgacaac 240
 gcctgcctgc tggccgagga gatcgtcacc gcgcgcggcg agcgggcgcg ctatttcggc 300
 agcgagggca agtcgtcggc cgagctggac aacctgccgt ccgacatctt ctacgcctg 360
 ctcaaccaac ccgagcgct gcaagccttc ttggcaggcg agacgcccg ggtacgccag 420
 ttgctcgaag gctacgccg cggtttcaac cgttctctcc gcgaggccga cggcaagacc 480
 accagttgcc ttggccagcc ctggctgcgg gccatcgca cccgatgacct gctgcgcctg 540
 acccgcgcc tgcctggcga agggggggtc ggccagttcg ccgacgcgt ggtggccgcc 600
 gcgcgcggc gagcggagaa ggtgccttg agcggcgagc aggcgttcca ggtcgccgag 660
 cagcgcgcc agcgttccg cctggagcgc ggcagcaacg ccattgccgt tggcagcgaa 720
 cgttcggcg acggcaagg catgctcctg gccaacccgc acttcccctg gaacggcgcg 780
 atgcgtttct accagatgca cctgaccatt cccggccggc tcgacgtgat gggggcctcg 840
 ctgcccggcc tgccgggtgt caacatcgcc ttacgcgcgc acctggcctg gacccacacg 900
 gtggatacct ccagccactt caccctgtat cgcctggcgc tggaccgaa ggaccgcgg 960
 cgctacctgg tcgacggctg ttgcgtccg ctggaggaga agtccgtcgc gatcgaggtg 1020
 cgcggcgccg acggcaagct gtcgcgcgtc gagcacaagg tctaccagtc gatctacggc 1080
 ccgctggtgg tctggcccg caagctggac tggaaaccga gcgaggccta tgcgtgcgt 1140
 gacgccaacc tggagaacac cggggtactg caacagtgtt actcgatcaa ccaggccagc 1200
 gacgtgcgg acctgcgcgc gcgcgtcgag gcgtacagg ggatcccctg ggtcaacacc 1260
 ctggccgcgg acgagcagg caacgcctg tacatgaacc agtcggtggt gccctacctg 1320
 aagccggaac tgattccgc ctgcgccatt ccgcaactgg tcgccgaagg cctgcggcc 1380
 ctccaggggc aggacagccg ctgtgcctgg agtcgcgacc cggccgcggc ccaggctggc 1440
 atcaccccg cggcgcaact gccgtgctg ttgcgcagg acttcgtgca gaactccaac 1500
 gacagcgct ggctgacca cccggcgagc ccgctgcagg gcttctcgcc cctggtcagc 1560
 caggagagc ccatcggtcc gcggcgcgcc tacgcctga gccggctaca gggcaagcag 1620
 ccgctggagg cgaagacgct cgaggagatg gtcaccgcca accatgtctt cagcgccgac 1680
 caggtgctgc cggacctgct ccgctgtgc cgcgacaacc agggcgagaa gtcccttgcc 1740
 cgcgcctgc cggccctggc gcagtgggac cgtggcgcca acctcgacag cggcagcgcc 1800
 ttctgtctact tccagcgctt catgcaacgc ttgcgcgaac tcgacggcg gcggaaggaa 1860
 ccgttcgatg cgcaacgtcc cctggatacg ccgcaaggca tcgccctcga ccggccgcag 1920
 gtggcgacc aggctgcgca ggcgtggcg gacgcggcg cggaggtgga gaagagcggg 1980
 attcccgagc gcgcgcgtg gggcgacctg caagtgcgca cccgtggcca ggaacgcac 2040
 gcgattcccg gcggcgatgg ccatttcggg gtctacaacg cgatccagag cgtccgcaag 2100
 ggcgaccacc tggaggtggc cggcggaact agctacatcc agctggtgac cttccccgag 2160
 gaagggccca aggtctcgcg gttgctggct ttctcccagt ccagcgatcc gcgctcgccg 2220
 cactaccgag accagaccga gctgttttcc cgccagcaat ggcagacctt gccgttcagc 2280
 gacaggcaga tcgacgcga cccgcaactg caacggctaa gcattcgca atga 2334

<210> 35

<211> 6390

<212> DNA

<213> Pseudomonas aeruginosa

<400> 35

gtgcgagggg tagccatgag tgcgtcagaa gacctgcaat ccgctgtgca accggccgcg 60
 agcgaagcgc tcgaaggatt cccgctgtct cccttgcaaga cccgcgcctg gcgcgcgcat 120
 gccgagcggc cggaaaatac ggttgctggc gtgcgcctgc acgccccggc cgatcccgtg 180
 gcgacgctgg agcggctgcg ccgggcgctg gacggcgagg cgcaactgcg cgtggcctac 240
 cggacgatgc cgggcatgag cctgcgggtg caggtactgg atgggcgcgc ggccgatctg 300
 ctggctcgagc gcctgcggg agacggcgac tgggcggac gcttcgcgcg cgaaagcgcg 360
 cgtctcgccg ctctgcgccct gggcggggaa ggccagccgg tactggcgct cggcctgctg 420
 ctggacggcg ccggagagac gctccagggg ctgttgctgg cggcgccggc gttcgtcgtc 480
 gatgcggcca gcctggtggc gctgctgocg cgcggcctgg ggccggccgg ccaggcgagc 540
 gcggacgagg gagacgagg gctgctgttc cagcatttct ccgagtgggc caacgaggcg 600
 ctggccggcg aagacggcga aagcgccagc ggttactggc gagagcaggc ggccgttgcg 660
 gcggagagtc cgtggcgct ggccggacgac ctgggcgaag gcgagtggac ggcgcgccgc 720
 ctgctgccgc gcgcgctgct cgaacgcctg gccgccaacg gcttgccgga ggccggccgc 780
 ctgctggcct ggaccaggt cgcggggcag ttccaggggc acgaggcct cccgctggaa 840

2025-06-06 10:00:00

[illegible][illegible]

gtgcaggaca tcgccgggct gaagatgcgc acccgcgccg gcatgggcga ctgccagggg 1260
 cggatgtgca tcggctactg cagcgatcgc ctgcgccgtg ccaccggacg ccacgacgtc 1320
 ggctggctgc ggccgcgttt cccgatcgat ccgatcccgt tttccgcatt ccagaacctc 1380
 ggtacggaag cctga 1395

<210> 37

<211> 801

<212> DNA

<213> Pseudomonas aeruginosa

<400> 37

atggcgctccg cccctaagga agaggagata aatatgattt attacttgat cggagtggcg 60
 ctattcatct tcatgctgga acagttgggt cccggctgga aattgcccaa ggtgagcacc 120
 tgggtggccc gggatgatctt cctcaacatc gtccagggtg cgatcgccct gctcgccggc 180
 atcacctgga acaaattgat gatggggcac agcctgctgc acacctcgga tgccttgcca 240
 ccactgctgg ccggcttcgc cgcctacttc gtcaaacacct tcgtcaccta ctggtggcat 300
 cgcgcgcgcc acgccaacga cagctctgg cggctgttcc accagttgca ccacgcgccg 360
 caacgcatcg aggtattcac ctcccttctac aagcatccga ccgagatggt cttcaactcg 420
 ctgctgggca gcttcgtcgc ctacgtgggt atgggcatca gcatcgaggc cggcgcctac 480
 tacatcatgt tcgccgcgct cggcgagatg ttctaccact cgaacctgcg caccgcgcac 540
 gtccctgggt acctgttcca gcgcccgag atgcaccgca tccaccacca gcgcgaccgt 600
 cacgagtgca actacagcga cttcccgatc tgggacatgt tggtcggcac ctacgagaac 660
 cccgcgcgca tcgacgagcc gcagggttc gccggcgaca aggagcagca gttcgtcgac 720
 atgctgctgt tccgcgacgt gcacagcctc cccggaaaaa cccagcccg cccgtcctg 780
 gtcaagcccc acgtcagggt a 801

<210> 38

<211> 20

<212> DNA

<213> Pseudomonas aeruginosa

<400> 38

acctgccccg aagggcaggt 20

<210> 39

<211> 468

<212> DNA

<213> Pseudomonas aeruginosa

<400> 39

gggttacctg gcacctacca gatcgtgtag ttgagccggt acgagcgctt tgtgttttat 60
 gcaatccaca tcagcgacca gggatgctgg ctatttgaaa cacttcacgg aatgacgctg 120
 aaagtcttcg cgacctcgtc tgtcgacact taacgaaagc attgcgaatc cattaccgac 180
 aggttttcaa aagaaaccgg ggatgaaact cctattgcct ttcgaaaatt ggaaacgaca 240
 ggcgaaacata tgtaacgcga aatttcaccc tacgtataaa caatgcgccc agcgaatatc 300
 gctcccttac cgagcgacga actcctgcgc gccagcgaat aaccgatgcc gcgagggaaa 360
 agtttctccg gcatacctgg agagccctct cggaggcggc gcatgaacgg tcagcggtac 420
 agggaaacac ccctcgacat cgagccgtct gcggcgcctt ctgagaca 468